



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C12Q</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/75358</b> <b>(43) International Publication Date:</b> 14 December 2000 (14.12.2000)
<b>(21) International Application Number:</b> PCT/US00/15364 <b>(22) International Filing Date:</b> 02 June 2000 (02.06.2000) <b>(30) Priority Data:</b> 60/137,659 04 June 1999 (04.06.1999) US <b>(60) Parent Application or Grant</b> MILLENNIUM PHARMACEUTICALS, INC. [/]; O. KHODADOUST, Mehran [/]; O. MANDRAGOURAS, Amy, E.; O.		<b>Published</b>
<b>(54) Title: NOVEL TOLL MOLECULES AND USES THEREFOR</b> <b>(54) Titre: MOLECULES TOLL ET LEURS UTILISATIONS</b>  <b>(57) Abstract</b> <p>The invention provides isolated nucleic acid molecules, designated TOLL nucleic acid molecules, which encode novel TOLL family members. The invention also provides antisense nucleic acid molecules, recombinant expression vectors containing TOLL nucleic acid molecules, host cells into which the expression vectors have been introduced, and nonhuman transgenic animals in which a TOLL gene has been introduced or disrupted. The invention still further provides isolated TOLL proteins, fusion proteins, antigenic peptides and anti-TOLL antibodies. Diagnostic methods utilizing compositions of the invention are also provided.</p> <b>(57) Abrégé</b> <p>Cette invention concerne des molécules d'acides nucléiques isolées dites molécules d'acides nucléiques TOLL, qui codent pour des membres de la famille TOLL. L'invention porte également sur des molécules d'acides nucléiques antisens, des vecteurs d'expression recombinants renfermant des molécules d'acides nucléiques TOLL, des cellules hôtes dans lesquels on a introduit des vecteurs d'expression et des animaux transgéniques non humains dans lesquels un gène TOLL a été introduit ou disloqué. Un autre aspect de l'invention concerne des protéines TOLL isolées, des protéines hybrides, des peptides antigéniques et des anticorps anti-TOLL. Sont également décrites des méthodes faisant intervenir des compositions selon l'invention.</p>		

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
14 December 2000 (14.12.2000)

PCT

(10) International Publication Number  
**WO 00/75358 A2**

- (51) International Patent Classification<sup>7</sup>: **C12Q** (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (21) International Application Number: PCT/US00/15364
- (22) International Filing Date: 2 June 2000 (02.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (30) Priority Data:  
60/137,659 4 June 1999 (04.06.1999) US
- (71) Applicant: MILLENNIUM PHARMACEUTICALS, INC. [US/US]; 75 Sidney Street, Cambridge, MA 02139 (US).
- (72) Inventor: KHODADOUST, Mehran; 106 Sewall Avenue #6, Brookline, MA 02445 (US).
- (74) Agents: MANDRAGOURAS, Amy, E.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 et al. (US).

**Published:**

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



**WO 00/75358 A2**

(54) Title: NOVEL TOLL MOLECULES AND USES THEREFOR

(57) Abstract: The invention provides isolated nucleic acid molecules, designated TOLL nucleic acid molecules, which encode novel TOLL family members. The invention also provides antisense nucleic acid molecules, recombinant expression vectors containing TOLL nucleic acid molecules, host cells into which the expression vectors have been introduced, and nonhuman transgenic animals in which a TOLL gene has been introduced or disrupted. The invention still further provides isolated TOLL proteins, fusion proteins, antigenic peptides and anti-TOLL antibodies. Diagnostic methods utilizing compositions of the invention are also provided.

**Description**

5

10

15

20

25

30

35

40

45

50

55

5

- 1 -

## NOVEL TOLL MOLECULES AND USES THEREFOR

10

Background of the Invention

The Toll protein in *Drosophila* is an integral membrane receptor involved in many different cellular processes ranging from cellular adhesion (Keith, F.J. et al., (1990) *EMBO J.* 9:4299-4306) to muscle formation (Halfon, M.S. et al., (1998) *Dev. Biol.* 199:164-174) to dorsal-ventral patterning during development (Hashimoto, C. et al., (1988) *Cell* 52:269-279). Studies in various other organisms have identified a number of related proteins, including the *Drosophila* 18-wheeler (Eldon, E., et al., (1994) *Development* 120:885-899), MstProx (Hashimoto, C. et al., (1988) *Cell* 52:269-279), and STSDm2245 proteins (Mitcham, J.L., et al., (1996) *J. Biol. Chem.* 271:5777-5783), the tobacco N gene product (Whitham, S. et al., (1994) *Cell* 78:1101-1115), mammalian and avian type-1 interleukin-1 receptors (Yamagata, M., et al. (1994) *Gene* 139:223-8), and human Toll-like proteins 1-5 (Rock, F.L., et al., (1998) *Proc. Natl. Acad. Sci. U.S.A.* 95:588-593).

30

Family members contain one or more specific conserved domains found in Toll. These include an extracellular leucine-rich repeat region, a domain containing a  $\beta/\alpha$ -class fold with  $\alpha$ -helices on both faces of a parallel  $\beta$  sheet, and a transmembrane domain (Hashimoto, C. et al., (1988) *Cell* 52:269-279; Rock, F.L., et al., (1998) *Proc. Natl. Acad. Sci. U.S.A.* 95:588-593). The specific functions of each of these domains within Toll have been explored. There is evidence that the extracellular leucine rich repeat region may promote cell to cell contact and adhesion due to the tendency of this domain to form an amphipathic structure with a predominantly apolar and charged surface (Gay, N.J., et al., (1991) *FEBS Lett* 291:87-91). The transmembrane domain serves to anchor the receptor into the membrane, and the cytosolic  $\beta/\alpha$ -class fold domain is posited to be a conformational trigger for signaling (Rock, F.L., et al., (1998) *Proc. Natl. Acad. Sci. U.S.A.* 95:588-593).

45

The conserved cytoplasmic domain between Toll and the avian/mammalian type 1 IL-1 receptor, containing the  $\beta/\alpha$ -class fold, suggested another potential function for Toll. The IL-1 receptor is involved in activation of the transcription factor NF $\kappa$ B in

50

55



5 response to stimulation by IL-1 in an innate immune response (O'Neill, L.A. et al.,  
(1998) *J. Leukoc. Biol.* 63: 650-657). It was found that mutation of certain residues in  
10 this domain not only abrogated the signaling activity of the type 1 IL-1 receptor, but also  
similarly impaired Toll function (Heguy, A. et al., (1992) *J. Biol. Chem.* 267:2606-  
5 2609). Further studies determined that Toll controls an immune response to fungal  
infection in *Drosophila* (Lemaitre, B., et al., (1996) *Cell* 86:973-983) and that 18-  
15 wheeler is critical for the normal functioning of the antibacterial response in these  
organisms (Williams, M.J., et al., (1997) *EMBO J.* 16:6120-6130). Similarly, studies of  
the tobacco N protein have demonstrated that this protein mediates resistance to tobacco  
20 mosaic virus (Whitham, S. et al., (1994) *Cell* 78:1101-1115). Thus, Toll family  
members may play key roles in host resistance to infection or in the activation of the  
immune response in general.

#### 25 Summary of the Invention

15 The present invention is based, at least in part, on the discovery of novel TOLL  
family members, referred to herein as "TOLL" nucleic acid and protein molecules. The  
TOLL molecules of the present invention are useful as targets for developing  
30 modulating agents to regulate a variety of cellular processes. Accordingly, in one  
aspect, this invention provides isolated nucleic acid molecules encoding TOLL proteins  
or biologically active portions thereof, as well as nucleic acid fragments suitable as  
20 primers or hybridization probes for the detection of TOLL-encoding nucleic acids.

35 In one embodiment, a TOLL nucleic acid molecule of the invention is at least  
50%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or more identical to the  
nucleotide sequence (e.g., to the entire length of the nucleotide sequence) shown in SEQ  
40 ID NO:1, 3, or 4 or the nucleotide sequence of the DNA insert of the plasmid deposited  
25 with ATCC as Accession Number \_\_\_\_\_, or a complement thereof.

45 In a preferred embodiment, the isolated nucleic acid molecule includes the  
nucleotide sequence shown SEQ ID NO:1, 3, or 4, or a complement thereof. In another  
embodiment, the nucleic acid molecule includes SEQ ID NO:3 and nucleotides 1-91 of  
30 SEQ ID NO:1. In another embodiment, the nucleic acid molecule includes SEQ ID  
50 NO:3 and nucleotides 1736-2147 of SEQ ID NO:1. In another preferred embodiment,

5 the nucleic acid molecule consists of the nucleotide sequence shown in SEQ ID NO:1, 3,  
or 4. In another preferred embodiment, the nucleic acid molecule includes a fragment of  
10 at least 491 nucleotides (e.g., 491 contiguous nucleotides) of the nucleotide sequence of  
SEQ ID NO:1 or 3, or a complement thereof.

5 In another embodiment, a TOLL nucleic acid molecule includes a nucleotide  
sequence encoding a protein having an amino acid sequence sufficiently homologous to  
15 the amino acid sequence of SEQ ID NO:2 or an amino acid sequence encoded by the  
DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_. In a  
preferred embodiment, a TOLL nucleic acid molecule includes a nucleotide sequence  
10 encoding a protein having an amino acid sequence at least 50%, 55%, 60%, 65%, 70%,  
20 75%, 80%, 85%, 90%, 95%, 98% or more homologous to the entire length of the amino  
acid sequence of SEQ ID NO:2 or the amino acid sequence encoded by the DNA insert  
of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_.

25 In another preferred embodiment, an isolated nucleic acid molecule encodes the  
15 amino acid sequence of human TOLL. In yet another preferred embodiment, the nucleic  
acid molecule includes a nucleotide sequence encoding a protein having the amino acid  
sequence of SEQ ID NO:2 or the amino acid sequence encoded by the DNA insert of the  
30 plasmid deposited with ATCC as Accession Number \_\_\_\_\_. In yet another preferred  
embodiment, the nucleic acid molecule is at least 491 nucleotides in length. In a further  
20 preferred embodiment, the nucleic acid molecule is at least 491 nucleotides in length and  
35 encodes a protein having a TOLL activity (as described herein).

Another embodiment of the invention features nucleic acid molecules, preferably  
TOLL nucleic acid molecules, which specifically detect TOLL nucleic acid molecules  
40 relative to nucleic acid molecules encoding non-TOLL proteins. For example, in one  
25 embodiment, such a nucleic acid molecule is at least 491, 491-500, 500-600, 600-700,  
700-800, 800-900, 900-1000, 1000-1100, 1100-1200, 1200-1300, 1300-1400, or 1400-  
45 1500 or more nucleotides in length and hybridizes under stringent conditions to a  
nucleic acid molecule comprising the nucleotide sequence shown in SEQ ID NO:1, the  
nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as  
30 Accession Number \_\_\_\_\_, or a complement thereof.

5 In preferred embodiments, the nucleic acid molecules are at least 15 (e.g.,  
contiguous) nucleotides in length and hybridize under stringent conditions to nucleotides  
1-547, or 1039-1322 of SEQ ID NO:1, or to nucleotides 1-763 of SEQ ID NO: 4. In  
10 other preferred embodiments, the nucleic acid molecules comprise nucleotides 1-547, or  
5 1039-1322 of SEQ ID NO:1, or nucleotides 1-763 of SEQ ID NO: 4.

In other preferred embodiments, the nucleic acid molecule encodes a naturally  
15 occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID  
NO:2 or an amino acid sequence encoded by the DNA insert of the plasmid deposited  
with ATCC as Accession Number \_\_\_\_\_, wherein the nucleic acid molecule hybridizes  
20 to a nucleic acid molecule comprising SEQ ID NO:1, 3, or 4 under stringent conditions.

Another embodiment of the invention provides an isolated nucleic acid molecule  
which is antisense to a TOLL nucleic acid molecule, e.g., the coding strand of a TOLL  
nucleic acid molecule.

25 Another aspect of the invention provides a vector comprising a TOLL nucleic  
15 acid molecule. In certain embodiments, the vector is a recombinant expression vector.  
In another embodiment, the invention provides a host cell containing a vector of the  
invention. In yet another embodiment, the invention provides a host cell containing a  
30 nucleic acid molecule of the invention. The invention also provides a method for  
producing a protein, preferably a TOLL protein, by culturing in a suitable medium, a  
20 host cell, e.g., a mammalian host cell such as a non-human mammalian cell, of the  
invention containing a recombinant expression vector, such that the protein is produced.

Another aspect of this invention features isolated or recombinant TOLL proteins  
and polypeptides. In one embodiment, the isolated protein, preferably a TOLL protein,  
40 includes at least one transmembrane domain. In another embodiment, the isolated  
25 protein, preferably a TOLL protein, includes a leucine-rich repeat (LRR) domain. In yet  
another embodiment, the isolated protein, preferably a TOLL protein, includes a domain  
containing a  $\beta/\alpha$ -class fold with  $\alpha$ -helices on both faces of a parallel  $\beta$  sheet. In a  
45 preferred embodiment, the protein, preferably a TOLL protein, includes at least one  
transmembrane domain and has an amino acid sequence at least about 50%, 55%, 60%,  
30 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or more homologous to the amino acid  
sequence of SEQ ID NO:2 or the amino acid sequence encoded by the DNA insert of the

5 plasmid deposited with ATCC as Accession Number \_\_\_\_\_. In another preferred  
embodiment, the protein, preferably a TOLL protein, includes a leucine-rich repeat  
10 (LRR) domain and has an amino acid sequence at least about 50%, 55%, 60%, 65%,  
70%, 75%, 80%, 85%, 90%, 95%, 98% or more homologous to the amino acid sequence  
5 of SEQ ID NO:2 or the amino acid sequence encoded by the DNA insert of the plasmid  
deposited with ATCC as Accession Number \_\_\_\_\_. In another preferred embodiment,  
15 the protein, preferably a TOLL protein, includes a domain containing a  $\beta/\alpha$ -class fold  
with  $\alpha$ -helices on both faces of a parallel  $\beta$  sheet and has an amino acid sequence at least  
about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or more  
20 homologous to the amino acid sequence of SEQ ID NO:2 or the amino acid sequence  
encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number  
\_\_\_\_\_. In yet another preferred embodiment, the protein, preferably a TOLL protein,  
includes at least one transmembrane domain, at least one leucine-rich repeat (LRR)  
25 domain and has an amino acid sequence at least about 50%, 55%, 60%, 65%, 70%, 75%,  
80%, 85%, 90%, 95%, 98% or more homologous to the amino acid sequence of SEQ ID  
15 NO:2 or the amino acid sequence encoded by the DNA insert of the plasmid deposited  
with ATCC as Accession Number \_\_\_\_\_. In yet another preferred embodiment, the  
30 protein, preferably a TOLL protein, includes at least one transmembrane domain and is  
encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes  
20 under stringent hybridization conditions to a nucleic acid molecule comprising the  
35 nucleotide sequence of SEQ ID NO:1, 3, or 4. In a further embodiment, the protein,  
preferably a TOLL protein, includes a leucine-rich repeat domain and is encoded by a  
nucleic acid molecule having a nucleotide sequence which hybridizes under stringent  
40 hybridization conditions to a nucleic acid molecule comprising the nucleotide sequence  
25 of SEQ ID NO:1, 3, or 4. In yet another preferred embodiment, the protein, preferably a  
TOLL protein, includes at least one domain containing a  $\beta/\alpha$ -class fold with  $\alpha$ -helices  
45 on both faces of a parallel  $\beta$  sheet and is encoded by a nucleic acid molecule having a  
nucleotide sequence which hybridizes under stringent hybridization conditions to a  
nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 1, 3, or 4. In  
30 yet another preferred embodiment, the protein, preferably a TOLL protein, includes at

5 least one transmembrane domain, at least one leucine-rich repeat (LRR) domain and is  
encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes  
under stringent hybridization conditions to a nucleic acid molecule comprising the  
10 nucleotide sequence of SEQ ID NO: 1, 3, or 4.

5 In another embodiment, the invention features fragments of the protein having  
the amino acid sequence of SEQ ID NO:2, wherein the fragment comprises at least 15  
15 amino acids (e.g., contiguous amino acids) of the amino acid sequence of SEQ ID NO:2  
or an amino acid sequence encoded by the DNA insert of the plasmid deposited with the  
ATCC as Accession Number \_\_\_\_\_. In another embodiment, the protein, preferably a  
20 TOLL protein, has the amino acid sequence of SEQ ID NO:2, respectively.

In another embodiment, the invention features an isolated protein, preferably a  
TOLL protein, which is encoded by a nucleic acid molecule consisting of a nucleotide  
sequence at least about 50%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or  
25 more homologous to a nucleotide sequence of SEQ ID NO:1, 3 or 4, or a complement  
15 thereof. This invention further features an isolated protein, preferably a TOLL protein,  
which is encoded by a nucleic acid molecule consisting of a nucleotide sequence which  
hybridizes under stringent hybridization conditions to a nucleic acid molecule  
30 comprising the nucleotide sequence of SEQ ID NO:1, 3, or 4, or a complement thereof.

The proteins of the present invention or portions thereof, e.g., biologically active  
20 portions thereof, can be operatively linked to a non-TOLL polypeptide (e.g.,  
35 heterologous amino acid sequences) to form fusion proteins. The invention further  
features antibodies, such as monoclonal or polyclonal antibodies, that specifically bind  
proteins of the invention, preferably TOLL proteins. In addition, the TOLL proteins or  
40 biologically active portions thereof can be incorporated into pharmaceutical  
25 compositions, which optionally include pharmaceutically acceptable carriers.

In another aspect, the present invention provides a method for detecting the  
presence of a TOLL nucleic acid molecule, protein or polypeptide in a biological sample  
45 by contacting the biological sample with an agent capable of detecting a TOLL nucleic  
acid molecule, protein or polypeptide such that the presence of a TOLL nucleic acid  
30 molecule, protein or polypeptide is detected in the biological sample.

5 In another aspect, the present invention provides a method for detecting the presence of TOLL activity in a biological sample by contacting the biological sample with an agent capable of detecting an indicator of TOLL activity such that the presence  
10 of TOLL activity is detected in the biological sample.

5 In another aspect, the invention provides a method for modulating TOLL activity comprising contacting a cell capable of expressing TOLL with an agent that modulates TOLL activity such that TOLL activity in the cell is modulated. In one embodiment, the  
15 agent inhibits TOLL activity. In another embodiment, the agent stimulates TOLL activity. In one embodiment, the agent is an antibody that specifically binds to a TOLL protein. In another embodiment, the agent modulates expression of TOLL by  
20 modulating transcription of a TOLL gene or translation of a TOLL mRNA. In yet another embodiment, the agent is a nucleic acid molecule having a nucleotide sequence that is antisense to the coding strand of a TOLL mRNA or a TOLL gene.

25 In one embodiment, the methods of the present invention are used to treat a subject having a disorder characterized by aberrant or unwanted TOLL protein or nucleic acid expression or activity by administering an agent which is a TOLL modulator to the subject. In one embodiment, the TOLL modulator is a TOLL protein.  
30 In another embodiment the TOLL modulator is a TOLL nucleic acid molecule. In yet another embodiment, the TOLL modulator is a peptide, peptidomimetic, or other small molecule.

35 The present invention also provides a diagnostic assay for identifying the presence or absence of a genetic alteration characterized by at least one of (i) aberrant modification or mutation of a gene encoding a TOLL protein; (ii) mis-regulation of the gene; and (iii) aberrant post-translational modification of a TOLL protein, wherein a  
40 wild-type form of the gene encodes a protein with a TOLL activity.

45 In another aspect the invention provides a method for identifying a compound that binds to or modulates the activity of a TOLL protein, by providing an indicator composition comprising a TOLL protein having TOLL activity, contacting the indicator composition with a test compound, and determining the effect of the test compound on  
50 TOLL activity in the indicator composition to identify a compound that modulates the activity of a TOLL protein.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

#### **Brief Description of the Drawings**

*Figure 1* depicts the cDNA sequence and predicted amino acid sequence of human TOLL. The nucleotide sequence corresponds to nucleic acids 1 to 2147 of SEQ ID NO:1. The amino acid sequence corresponds to amino acids 1 to 548 of SEQ ID NO:

2. The coding region without the 5' and 3' untranslated regions of the human TOLL gene is shown in SEQ ID NO:3.

*Figure 2* depicts the results of a search which was performed against the HMM database and which resulted in the identification of a leucine-rich repeat domain in the human TOLL protein.

*Figure 3* depicts the nucleotide sequence of mouse TOLL. The nucleotide sequence corresponds to nucleic acids 1 to 763 of SEQ ID NO:4.

#### **Detailed Description of the Invention**

The present invention is based, at least in part, on the discovery of novel nucleic acid and protein molecules which may be Toll family members, referred to herein as "TOLL" nucleic acid and protein molecules. Due to their homology with Toll family members, the TOLL molecules of the present invention may be membrane proteins, e.g., membrane proteins which function as receptors, and they may be involved in immune signaling mechanisms.

As used herein, the term "immune signaling mechanisms" includes the cellular mechanisms involved in the development and regulation of an immune response, e.g., an immune response to the presence of a pathogen or antigen in a subject, e.g., a mammal such as a human. In mammals, the initial detection of a "foreign" pathogen or antigen results in the triggering of the innate immune response, in which specialized cells are recruited to engulf and destroy the invader, while chemical signals are simultaneously generated and released to stimulate the activation of the acquired immune system. The TOLL molecules of the present invention may be involved in pathways involved in the

5 initial detection of these invaders, and/or in the recruitment of specialized cells such as macrophages, and/or in stimulation of cytokine production and release.

10 Thus, the TOLL molecules, by participating in immune signaling mechanisms, may be useful in the development of novel diagnostic targets and therapeutic agents to  
5 regulate immune and inflammatory responses in a variety of disorders, diseases, or conditions which are characterized by a deregulated, e.g., upregulated or downregulated, immune or inflammatory response. For example, the TOLL molecules may provide  
15 novel diagnostic targets and therapeutic agents for controlling disorders, diseases, or conditions related to misregulation of an immune response, such as rheumatoid arthritis, systemic lupus erythematosus, myasthenia gravis, Grave's disease, Sjogren syndrome,  
20 polymyositis and dermatomyositis, psoriasis, pemphigus vulgaris, bullous pemphigoid, inflammatory bowel disease, Kawasaki disease, asthma, and graft vs. host disease.

The term "family" when referring to the protein and nucleic acid molecules of  
25 the invention is intended to mean two or more proteins or nucleic acid molecules having  
15 a common structural domain or motif and having sufficient amino acid or nucleotide sequence homology as defined herein. Such family members can be naturally or non-naturally occurring and can be from either the same or different species. For example, a  
30 family can contain a first protein of human origin, as well as other, distinct proteins of human origin or alternatively, can contain homologues of non-human origin. Members  
20 of a family may also have common functional characteristics.

35 For example, the family of TOLL proteins include at least one "transmembrane domain". As used herein, the term "transmembrane domain" includes an amino acid sequence of about 15 amino acid residues in length which spans the plasma membrane. More preferably, a transmembrane domain includes about at least 20, 25, 30, 35, 40, or  
40 45 amino acid residues and spans the plasma membrane. Transmembrane domains are rich in hydrophobic residues, and typically have an  $\alpha$ -helical structure. In a preferred embodiment, at least 50%, 60%, 70%, 80%, 90%, 95% or more of the amino acids of a  
45 transmembrane domain are hydrophobic, e.g., leucines, isoleucines, tyrosines, or tryptophans. Transmembrane domains are described in, for example, Zagotta W.N. et  
30 al, (1996) *Annual Rev. Neurosci.* 19: 235-63, the contents of which are incorporated



5           herein by reference. Amino acid residues 456-480 of the human TOLL protein comprise a transmembrane domain.

10           In another embodiment, a TOLL of the present invention is identified based on the presence of a "leucine-rich repeat" domain in the protein or corresponding nucleic acid molecule. As used herein, the term "leucine-rich repeat" domain includes a protein domain having an amino acid sequence of about 50 amino acid residues and having a bit score for the alignment of the sequence to the leucine-rich repeat domain (LRR) of at least about 15. Preferably, a leucine-rich repeat domain includes at least about 20-60, more preferably about 20-50 amino acid residues, or about 20-40 amino acids and has a bit score for the alignment of the sequence to the leucine-rich repeat domain (LRR) of at least 20, 30, 40, 50 or greater. The leucine-rich repeat domain (LRR) has been assigned the PFAM Accession PF00560 (<http://genome.wustl.edu/Pfam/html>). Leucine-rich repeat domains are described in, for example, Kobe, B. et al., (1994) *Trends in Biol. Sci.* 19:415-421, the contents of which are incorporated herein by reference.

15           To identify the presence of a leucine-rich repeat domain in a TOLL protein, and make the determination that a protein of interest has a particular profile, the amino acid sequence of the protein is searched against a database of HMMs (e.g., the Pfam database, release 2.1) using the default parameters ([http://www.sanger.ac.uk/Software/Pfam/HMM\\_search](http://www.sanger.ac.uk/Software/Pfam/HMM_search)). For example, the hmmsf program, which is available as part of the HMMER package of search programs, is a family specific default program for MILPAT0063 and a score of 15 is the default threshold score for determining a hit. Alternatively, the threshold score for determining a hit can be lowered (e.g., to 8 bits). A description of the Pfam database can be found in Sonhammer et al. (1997) *Proteins* 28(3):405-420 and a detailed description of HMMs can be found, for example, in Gribskov et al. (1990) *Meth. Enzymol.* 183:146-159; Gribskov et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:4355-4358; Krogh et al. (1994) *J. Mol. Biol.* 235:1501-1531; and Stultz et al. (1993) *Protein Sci.* 2:305-314, the contents of which are incorporated herein by reference. A search was performed against the HMM database resulting in the identification of a leucine-rich repeat domain in the amino acid sequence of human TOLL (SEQ ID NO:2) at about residues 58-105, 106-153, 154-201.

202-250, 251-297, and 298-345 of SEQ ID NO:2. The results of the search are set forth in Figure 2.

Accordingly, TOLL proteins having at least 50-60% homology, preferably about 60-70%, more preferably about 70-80%, or about 80-90% homology with a leucine-rich repeat domain of human TOLL (e.g., residues 58-105, 106-153, 154-201, 202-250, 251-297, or 298-345 of SEQ ID NO:2) or with a TM domain of human TOLL (e.g., residues 456-480 and 483-507 of SEQ ID NO:2) are within the scope of the invention.

In another embodiment, a TOLL molecule of the present invention is identified based on the presence of a  $\beta/\alpha$ -class fold with  $\alpha$ -helices on both faces of a parallel  $\beta$  sheet in the protein molecule. As used herein, the term " $\beta/\alpha$ -class fold with  $\alpha$ -helices on both faces of a parallel  $\beta$  sheet" includes a domain having a  $\beta$  sheet structure in which the longer and more hydrophobic  $\beta$  strands are predicted to form interior staves in the beta sheet, while the more amphipathic  $\beta$  strands are at the edge of the sheet, with  $\alpha$ -helices on both faces of the sheet (see Rock, F.L. *et al.* (1998) *Proc. Natl. Acad. Sci. U.S.A.* 95: 588-593, the contents of which are incorporated herein by reference). An  $\alpha$  helix is a rodlike coiled structure stabilized by hydrogen bonds between the NH and CO groups of the polypeptide chain, while a  $\beta$  sheet is a sheet comprised of different  $\beta$  strands which may be parallel or antiparallel, in which the strands are almost fully extended (as opposed to the tightly coiled  $\alpha$ -helix); such structures are known in the art.

Isolated proteins of the present invention, preferably TOLL proteins, have an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2 or are encoded by a nucleotide sequence sufficiently homologous to SEQ ID NO:1, 3, or 4. As used herein, the term "sufficiently homologous" refers to a first amino acid or nucleotide sequence which contains a sufficient or minimum number of identical or equivalent (e.g., an amino acid residue which has a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or nucleotide sequences share common structural domains or motifs and/or a common functional activity. For example, amino acid or nucleotide sequences which share common structural domains have at least 30%, 40%, or 50% homology, preferably 60% homology, more preferably 70%-80%, and even more preferably 90-95% homology across the amino

5 acid sequences of the domains and contain at least one and preferably two structural domains or motifs, are defined herein as sufficiently homologous. Furthermore, amino acid  
10 or nucleotide sequences which share at least 30%, 40%, or 50%, preferably 60%, more preferably 70-80%, or 90-95% homology and share a common functional activity are  
5 defined herein as sufficiently homologous.

As used interchangeably herein, an "TOLL activity", "biological activity of  
15 TOLL" or "functional activity of TOLL", refers to an activity exerted by a TOLL protein, polypeptide or nucleic acid molecule on a TOLL responsive cell or on a TOLL protein substrate, as determined *in vivo*, or *in vitro*, according to standard techniques. In  
20 one embodiment, a TOLL activity is a direct activity, such as an association with a TOLL-target molecule. As used herein, a "target molecule" or "binding partner" is a molecule with which a TOLL protein binds or interacts in nature, such that TOLL-mediated function is achieved. A TOLL target molecule can be a non-TOLL molecule  
25 or a TOLL protein or polypeptide of the present invention. In an exemplary  
15 embodiment, a TOLL target molecule is a TOLL ligand. Alternatively, a TOLL activity is an indirect activity, such as a cellular signaling activity mediated by interaction of the TOLL protein with a TOLL ligand.

Accordingly, another embodiment of the invention features isolated TOLL  
30 proteins and polypeptides having a TOLL activity. Preferred proteins are TOLL  
20 proteins having at least one transmembrane domain, and, preferably, a TOLL activity.  
35 Other preferred proteins are TOLL proteins having at least one leucine-rich repeat domain and, preferably, a TOLL activity. Other preferred proteins are TOLL proteins having at least one domain containing a  $\beta/\alpha$ -class fold with  $\alpha$ -helices on both faces of a  
40 parallel  $\beta$  sheet and, preferably, a TOLL activity. Additional preferred proteins have at  
25 least one transmembrane domain and/or a leucine-rich repeat domain, and are, preferably, encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule  
45 comprising the nucleotide sequence of SEQ ID NO:1, 3, or 4.

The nucleotide sequence of the isolated human TOLL cDNA and the predicted  
30 amino acid sequence of the human TOLL polypeptide are shown in Figure 1 and in SEQ ID NOs:1 and 2, respectively. The nucleotide sequence of the isolated partial mouse  
50

5 TOLL DNA is shown in Figure 3. A plasmid containing the nucleotide sequence  
encoding human TOLL was deposited with the American Type Culture Collection  
10 (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209, on \_\_\_\_\_ and  
assigned Accession Number \_\_\_\_\_. A plasmid containing the nucleotide sequence  
5 encoding partial mouse TOLL was deposited with the American Type Culture  
Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209, on \_\_\_\_\_  
15 and assigned Accession Number \_\_\_\_\_. These deposits will be maintained under the  
terms of the Budapest Treaty on the International Recognition of the Deposit of  
Microorganisms for the Purposes of Patent Procedure. These deposits were made  
10 merely as a convenience for those of skill in the art and is not an admission that a  
deposit is required under 35 U.S.C. §112.

20 The human TOLL gene, which is approximately 2147 nucleotides in length,  
encodes a protein having a molecular weight of approximately 75 kD and which is  
25 approximately 548 amino acid residues in length. The partial mouse TOLL nucleotide  
15 sequence is approximately 763 nucleotides in length.

30 Various aspects of the invention are described in further detail in the following  
subsections:

#### I. Isolated Nucleic Acid Molecules

20 One aspect of the invention pertains to isolated nucleic acid molecules that  
35 encode TOLL proteins or biologically active portions thereof, as well as nucleic acid  
fragments sufficient for use as hybridization probes to identify TOLL-encoding nucleic  
acid molecules (e.g., TOLL mRNA) and fragments for use as PCR primers for the  
40 amplification or mutation of TOLL nucleic acid molecules. As used herein, the term  
25 "nucleic acid molecule" is intended to include DNA molecules (e.g., cDNA or genomic  
DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated  
using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-  
45 stranded, but preferably is double-stranded DNA.

30 The term "isolated nucleic acid molecule" includes nucleic acid molecules which  
are separated from other nucleic acid molecules which are present in the natural source  
50 of the nucleic acid. For example, with regards to genomic DNA, the term "isolated"

5 includes nucleic acid molecules which are separated from the chromosome with which  
the genomic DNA is naturally associated. Preferably, an "isolated" nucleic acid is free  
10 of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and  
3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic  
5 acid is derived. For example, in various embodiments, the isolated TOLL nucleic acid  
molecule can contain less than about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb of  
15 nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA  
of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid  
molecule, such as a cDNA molecule, can be substantially free of other cellular material,  
20 or culture medium when produced by recombinant techniques, or substantially free of  
chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, e.g., a nucleic acid molecule  
having the nucleotide sequence of SEQ ID NO:1, 3, or 4, or the nucleotide sequence of  
25 the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_, or a  
15 portion thereof, can be isolated using standard molecular biology techniques and the  
sequence information provided herein. Using all or portion of the nucleic acid sequence  
of SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid  
30 deposited with ATCC as Accession Number \_\_\_\_\_, as a hybridization probe, TOLL  
nucleic acid molecules can be isolated using standard hybridization and cloning  
20 techniques (e.g., as described in Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular  
Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring  
35 Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

Moreover, a nucleic acid molecule encompassing all or a portion of SEQ ID  
40 NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited  
25 with ATCC as Accession Number \_\_\_\_\_ can be isolated by the polymerase chain  
reaction (PCR) using synthetic oligonucleotide primers designed based upon the  
sequence of SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the  
45 plasmid deposited with ATCC as Accession Number \_\_\_\_\_.

A nucleic acid of the invention can be amplified using cDNA, mRNA or  
30 alternatively, genomic DNA, as a template and appropriate oligonucleotide primers  
50 according to standard PCR amplification techniques. The nucleic acid so amplified can

5 be cloned into an appropriate vector and characterized by DNA sequence analysis.  
Furthermore, oligonucleotides corresponding to TOLL nucleotide sequences can be  
10 prepared by standard synthetic techniques, e.g., using an automated DNA synthesizer.

In a preferred embodiment, an isolated nucleic acid molecule of the invention  
5 comprises the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO: 4. The  
sequence of SEQ ID NO:1 corresponds to the human TOLL cDNA. This cDNA  
15 comprises sequences encoding the human TOLL protein (i.e., "the coding region", from  
nucleotides 92-1735), as well as 5' untranslated sequences (nucleotides 1-91) and 3'  
untranslated sequences (nucleotides 1736-2147). Alternatively, the nucleic acid  
20 molecule can comprise only the coding region of SEQ ID NO:1 (e.g., nucleotides 92-  
1735, corresponding to SEQ ID NO:3). The sequence of SEQ ID NO:4 corresponds to a  
fragment of the mouse TOLL cDNA.

In another preferred embodiment, an isolated nucleic acid molecule of the  
25 invention comprises a nucleic acid molecule which is a complement of the nucleotide  
15 sequence shown in SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert  
of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_, or a portion of any of  
these nucleotide sequences. A nucleic acid molecule which is complementary to the  
30 nucleotide sequence shown in SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the  
DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_, is one  
20 which is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1,  
3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited with  
35 ATCC as Accession Number \_\_\_\_\_, such that it can hybridize to the nucleotide  
sequence shown in SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert  
40 of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_, thereby forming a  
25 stable duplex.

In still another preferred embodiment, an isolated nucleic acid molecule of the  
45 present invention comprises a nucleotide sequence which is at least about 50%, 60%,  
65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or more homologous to the entire length of  
the nucleotide sequence shown in SEQ ID NO:1, 3, or 4, or the entire length of the  
30 nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as  
50 Accession Number \_\_\_\_\_, or a portion of any of these nucleotide sequences.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the nucleic acid sequence of SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_, for example, a fragment which can be used as a probe or primer or a fragment encoding a portion of a TOLL protein, e.g., a biologically active portion of a TOLL protein. The nucleotide sequence determined from the cloning of the TOLL gene allows for the generation of probes and primers designed for use in identifying and/or cloning other TOLL family members, as well as TOLL homologues from other species. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12 or 15, preferably about 20 or 25, more preferably about 30, 35, 40, 45, 50, 55, 60, 65, or 75 consecutive nucleotides of a sense sequence of SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_, of an anti-sense sequence of SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_, or of a naturally occurring allelic variant or mutant of SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_. In an exemplary embodiment, a nucleic acid molecule of the present invention comprises a nucleotide sequence which is greater than 491, 491-500, 500-550, 550-600, 600-650, 650-700, 700-750, 750-800, 800-850, 850-900, 900-950, 950-1000, or more nucleotides in length and hybridizes under stringent hybridization conditions to a nucleic acid molecule of SEQ ID NO:1 or 3, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_.

Probes based on the TOLL nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, e.g., the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress a TOLL protein, such as by measuring a level of a TOLL-

5 encoding nucleic acid in a sample of cells from a subject e.g., detecting TOLL mRNA levels or determining whether a genomic TOLL gene has been mutated or deleted.

10 A nucleic acid fragment encoding a "biologically active portion of a TOLL protein" can be prepared by isolating a portion of the nucleotide sequence of SEQ ID  
5 NO:1, 3, or 4 or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_, which encodes a polypeptide having a TOLL biological activity (the biological activities of the TOLL proteins are described herein),  
15 expressing the encoded portion of the TOLL protein (e.g., by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the TOLL protein.

20 The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence shown in SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_, due to degeneracy of the genetic code and thus encode the same TOLL proteins as those  
25 encoded by the nucleotide sequence shown in SEQ ID NO:1, 3, or 4, or the nucleotide  
15 sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ  
30 ID NO:2.

In addition to the TOLL nucleotide sequences shown in SEQ ID NO:1, 3, or 4, or  
20 the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as  
35 Accession Number \_\_\_\_\_, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of the TOLL proteins may exist within a population (e.g., the human population). Such genetic  
40 polymorphism in the TOLL genes may exist among individuals within a population due  
25 to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules which include an open reading frame encoding a TOLL protein, preferably a mammalian TOLL protein, and can further include non-coding  
45 regulatory sequences, and introns.

Allelic variants of human TOLL include both functional and non-functional  
30 TOLL proteins. Functional allelic variants are naturally occurring amino acid sequence variants of the human TOLL protein that maintain the ability to bind a TOLL ligand  
50



5 and/or to modulate signaling mechanisms involved in a TOLL related mechanism.  
Functional allelic variants will typically contain only conservative substitution of one or  
10 more amino acids of SEQ ID NO:2 or substitution, deletion or insertion of non-critical  
residues in non-critical regions of the protein.

5 Non-functional allelic variants are naturally occurring amino acid sequence  
variants of the human TOLL protein that do not have the ability to either bind a TOLL  
15 ligand and/or to modulate signaling mechanisms involved in, for example, an immune  
response. Non-functional allelic variants will typically contain a non-conservative  
substitution, a deletion, or insertion or premature truncation of the amino acid sequence  
20 of SEQ ID NO:2 or a substitution, insertion or deletion in critical residues or critical  
regions.

The present invention further provides non-human orthologues of the human  
TOLL protein. Orthologues of the human TOLL protein are proteins that are isolated  
25 from non-human organisms and possess the same TOLL ligand binding properties as the  
15 human TOLL protein. Orthologues of the human TOLL protein can readily be  
identified as comprising an amino acid sequence that is substantially homologous to  
SEQ ID NO:2.

30 Moreover, nucleic acid molecules encoding other TOLL family members and,  
thus, which have a nucleotide sequence which differs from the TOLL sequences of SEQ  
20 ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited  
35 with ATCC as Accession Number \_\_\_\_\_ are intended to be within the scope of the  
invention. For example, another TOLL cDNA can be identified based on the nucleotide  
sequence of human TOLL. Moreover, nucleic acid molecules encoding TOLL proteins  
40 from different species, and which, thus, have a nucleotide sequence which differs from  
25 the TOLL sequences of SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA  
insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_ are intended to  
be within the scope of the invention. For example, a mouse TOLL cDNA can be  
45 identified based on the nucleotide sequence of a human TOLL.

Nucleic acid molecules corresponding to natural allelic variants and homologues  
30 of the TOLL cDNAs of the invention can be isolated based on their homology to the  
50 TOLL nucleic acids disclosed herein using the cDNAs disclosed herein, or a portion

thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. Nucleic acid molecules corresponding to natural allelic variants and homologues of the TOLL cDNAs of the invention can further be isolated by mapping to the same chromosome or locus as the TOLL gene.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15, 20, 25, 30 or more nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_. In other embodiment, the nucleic acid is at least 30, 50, 100, 150, 200, 250, 253, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, or 950 nucleotides in length. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other. Preferably, the conditions are such that sequences at least about 70%, more preferably at least about 80%, even more preferably at least about 85% or 90% homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50°C, preferably at 55°C, more preferably at 60°C, and even more preferably at 65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1, 3, or 4 or corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (e.g., encodes a natural protein).

In addition to naturally-occurring allelic variants of the TOLL sequences that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequences of SEQ ID NO:1, 3, or 4, or the

5 nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession  
Number \_\_\_\_\_. thereby leading to changes in the amino acid sequence of the encoded  
TOLL proteins, without altering the functional ability of the TOLL proteins. For  
10 example, nucleotide substitutions leading to amino acid substitutions at "non-essential"  
5 amino acid residues can be made in the sequence of SEQ ID NO:1, 3, or 4, or the  
nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession  
Number \_\_\_\_\_. A "non-essential" amino acid residue is a residue that can be altered  
15 from the wild-type sequence of TOLL (e.g., the sequence of SEQ ID NO:2) without  
altering the biological activity, whereas an "essential" amino acid residue is required for  
10 biological activity. For example, amino acid residues that are conserved among the  
TOLL proteins of the present invention, e.g., those present in the transmembrane domain,  
are predicted to be particularly unamenable to alteration. Furthermore, additional amino  
20 acid residues that are conserved between the TOLL proteins of the present invention and  
other members of the TOLL family are not likely to be amenable to alteration.

15 Accordingly, another aspect of the invention pertains to nucleic acid molecules  
encoding TOLL proteins that contain changes in amino acid residues that are not  
essential for activity. Such TOLL proteins differ in amino acid sequence from SEQ ID  
30 NO:2, yet retain biological activity. In one embodiment, the isolated nucleic acid  
molecule comprises a nucleotide sequence encoding a protein, wherein the protein  
20 comprises an amino acid sequence at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%,  
35 85%, 90%, 95%, 98% or more homologous to SEQ ID NO:2.

An isolated nucleic acid molecule encoding a TOLL protein homologous to the  
protein of SEQ ID NO:2 can be created by introducing one or more nucleotide  
40 substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:1, 3, or  
25 4, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as  
Accession Number \_\_\_\_\_. such that one or more amino acid substitutions, additions or  
deletions are introduced into the encoded protein. Mutations can be introduced into SEQ  
45 ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid deposited  
with ATCC as Accession Number \_\_\_\_\_ by standard techniques, such as site-directed  
30 mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid  
substitutions are made at one or more predicted non-essential amino acid residues. A

5 "conservative amino acid substitution" is one in which the amino acid residue is replaced  
with an amino acid residue having a similar side chain. Families of amino acid residues  
having similar side chains have been defined in the art. These families include amino  
10 acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g.,  
5 aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine,  
glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine,  
15 valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-  
branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g.,  
tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid  
20 residue in a TOLL protein is preferably replaced with another amino acid residue from  
the same side chain family. Alternatively, in another embodiment, mutations can be  
introduced randomly along all or part of a TOLL coding sequence, such as by saturation  
mutagenesis, and the resultant mutants can be screened for TOLL biological activity to  
25 identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1, 3, or 4, or  
15 the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as  
Accession Number \_\_\_\_\_, the encoded protein can be expressed recombinantly and the  
activity of the protein can be determined.

30 In a preferred embodiment, a mutant TOLL protein can be assayed for the ability  
to (1) interact with a non-TOLL protein molecule, e.g., a TOLL ligand; (2) activate a  
20 TOLL-dependent signal transduction pathway; or (3) modulate signaling mechanisms  
involved in, for example, an immune response.

35 In addition to the nucleic acid molecules encoding TOLL proteins described  
above, another aspect of the invention pertains to isolated nucleic acid molecules which  
are antisense thereto. An "antisense" nucleic acid comprises a nucleotide sequence which  
40 is complementary to a "sense" nucleic acid encoding a protein, e.g., complementary to  
25 the coding strand of a double-stranded cDNA molecule or complementary to an mRNA  
sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic  
acid. The antisense nucleic acid can be complementary to an entire TOLL coding strand,  
45 or to only a portion thereof. In one embodiment, an antisense nucleic acid molecule is  
30 antisense to a "coding region" of the coding strand of a nucleotide sequence encoding  
TOLL. The term "coding region" refers to the region of the nucleotide sequence  
50

5 comprising codons which are translated into amino acid residues (e.g., the coding region of human TOLL corresponds to SEQ ID NO:3). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a  
10 nucleotide sequence encoding TOLL. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (i.e., also referred to as 5' and 3' untranslated regions).

15 Given the coding strand sequences encoding TOLL disclosed herein (e.g., SEQ ID NO:3), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be  
20 complementary to the entire coding region of TOLL mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of TOLL mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of TOLL mRNA. An antisense  
25 oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art.  
30 For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase  
35 the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-  
40 carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-  
45 methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutosine, pseudouracil, queosine, 2-

thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a TOLL protein to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention include direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an  $\alpha$ -anomeric nucleic acid molecule. An  $\alpha$ -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual  $\beta$ -units, the strands run parallel to each other (Gaultier et al. (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue et al. (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue et al. (1987) *FEBS Lett.* 215:327-330).

5 In still another embodiment, an antisense nucleic acid of the invention is a  
ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are  
capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they  
10 have a complementary region. Thus, ribozymes (e.g., hammerhead ribozymes  
5 (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to  
catalytically cleave TOLL mRNA transcripts to thereby inhibit translation of TOLL  
mRNA. A ribozyme having specificity for a TOLL-encoding nucleic acid can be  
15 designed based upon the nucleotide sequence of a TOLL cDNA disclosed herein (i.e.,  
SEQ ID NO:1, 3, or 4, or the nucleotide sequence of the DNA insert of the plasmid  
10 deposited with ATCC as Accession Number \_\_\_\_). For example, a derivative of a  
20 *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the  
active site is complementary to the nucleotide sequence to be cleaved in a TOLL-  
encoding mRNA. See, e.g., Cech et al. U.S. Patent No. 4,987,071; and Cech et al. U.S.  
25 Patent No. 5,116,742. Alternatively, TOLL mRNA can be used to select a catalytic  
15 RNA having a specific ribonuclease activity from a pool of RNA molecules. See, e.g.,  
Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

30 Alternatively, TOLL gene expression can be inhibited by targeting nucleotide  
sequences complementary to the regulatory region of the TOLL (e.g., the TOLL  
promoter and/or enhancers) to form triple helical structures that prevent transcription of  
20 the TOLL gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.*  
35 6(6):569-84; Helene, C. et al. (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L.J.  
(1992) *Bioassays* 14(12):807-15.

In yet another embodiment, the TOLL nucleic acid molecules of the present  
40 invention can be modified at the base moiety, sugar moiety or phosphate backbone to  
25 improve, e.g., the stability, hybridization, or solubility of the molecule. For example,  
the deoxyribose phosphate backbone of the nucleic acid molecules can be modified to  
generate peptide nucleic acids (see Hyrup B. et al. (1996) *Bioorganic & Medicinal*  
45 *Chemistry* 4 (1): 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs"  
refer to nucleic acid mimics, e.g., DNA mimics, in which the deoxyribose phosphate  
30 backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases  
are retained. The neutral backbone of PNAs has been shown to allow for specific

5 hybridization to DNA and RNA under conditions of low ionic strength. The synthesis  
of PNA oligomers can be performed using standard solid phase peptide synthesis  
protocols as described in Hyrup B. et al. (1996) *supra*; Perry-O'Keefe et al. Proc. Natl.  
10 Acad. Sci. 93: 14670-675.

5 PNAs of TOLL nucleic acid molecules can be used in therapeutic and diagnostic  
applications. For example, PNAs can be used as antisense or antigene agents for  
sequence-specific modulation of gene expression by, for example, inducing transcription  
15 or translation arrest or inhibiting replication. PNAs of TOLL nucleic acid molecules can  
also be used in the analysis of single base pair mutations in a gene, (e.g., by PNA-  
20 directed PCR clamping); as 'artificial restriction enzymes' when used in combination  
with other enzymes, (e.g., S1 nucleases (Hyrup B. (1996) *supra*)); or as probes or  
primers for DNA sequencing or hybridization (Hyrup B. et al. (1996) *supra*; Perry-  
O'Keefe *supra*).

25 In another embodiment, PNAs of TOLL can be modified, (e.g., to enhance their  
15 stability or cellular uptake), by attaching lipophilic or other helper groups to PNA, by  
the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of  
drug delivery known in the art. For example, PNA-DNA chimeras of TOLL nucleic  
30 acid molecules can be generated which may combine the advantageous properties of  
PNA and DNA. Such chimeras allow DNA recognition enzymes, (e.g., RNase H and  
20 DNA polymerases), to interact with the DNA portion while the PNA portion would  
provide high binding affinity and specificity. PNA-DNA chimeras can be linked using  
35 linkers of appropriate lengths selected in terms of base stacking, number of bonds  
between the nucleobases, and orientation (Hyrup B. (1996) *supra*). The synthesis of  
PNA-DNA chimeras can be performed as described in Hyrup B. (1996) *supra* and Finn  
40 P.J. et al. (1996) *Nucleic Acids Res.* 24 (17): 3357-63. For example, a DNA chain can  
be synthesized on a solid support using standard phosphoramidite coupling chemistry  
and modified nucleoside analogs, e.g., 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine  
45 phosphoramidite, can be used as a between the PNA and the 5' end of DNA (Mag, M. et  
al. (1989) *Nucleic Acid Res.* 17: 5973-88). PNA monomers are then coupled in a  
30 stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA  
segment (Finn P.J. et al. (1996) *supra*). Alternatively, chimeric molecules can be  
50



5 synthesized with a 5' DNA segment and a 3' PNA segment (Peteraser, K.H. et al. (1975) *Bioorganic Med. Chem. Lett.* 5: 1119-11124).

10 In other embodiments, the oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors *in vivo*), or agents facilitating  
5 transport across the cell membrane (see, e.g., Letsinger et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:6553-6556; Lemaitre et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, e.g., PCT Publication  
15 No. W089/10134). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (See, e.g., Krol et al. (1988) *Bio-Techniques* 6:958-976) or  
20 intercalating agents. (See, e.g., Zon (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, (e.g., a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent).

## 25 II. Isolated TOLL Proteins and Anti-TOLL Antibodies

15 One aspect of the invention pertains to isolated TOLL proteins, and biologically active portions thereof, as well as polypeptide fragments suitable for use as immunogens to raise anti-TOLL antibodies. In one embodiment, native TOLL proteins can be  
30 isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, TOLL proteins are  
20 produced by recombinant DNA techniques. Alternative to recombinant expression, a TOLL protein or polypeptide can be synthesized chemically using standard peptide  
35 synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is  
40 substantially free of cellular material or other contaminating proteins from the cell or  
25 tissue source from which the TOLL protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language  
"substantially free of cellular material" includes preparations of TOLL protein in which  
45 the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of  
30 cellular material" includes preparations of TOLL protein having less than about 30% (by  
50 dry weight) of non-TOLL protein (also referred to herein as a "contaminating protein"),

5 more preferably less than about 20% of non-TOLL protein, still more preferably less than about 10% of non-TOLL protein, and most preferably less than about 5% non-TOLL protein. When the TOLL protein or biologically active portion thereof is  
10 recombinantly produced, it is also preferably substantially free of culture medium, i.e.,  
5 culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

15 The language "substantially free of chemical precursors or other chemicals" includes preparations of TOLL protein in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. In one  
20 embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of TOLL protein having less than about 30% (by dry weight) of chemical precursors or non-TOLL chemicals, more preferably less than about 20%  
25 chemical precursors or non-TOLL chemicals, still more preferably less than about 10% chemical precursors or non-TOLL chemicals, and most preferably less than about 5%  
15 chemical precursors or non-TOLL chemicals.

As used herein, a "biologically active portion" of a TOLL protein includes a  
30 fragment of a TOLL protein which participates in an interaction between a TOLL molecule and a non-TOLL molecule. Biologically active portions of a TOLL protein include peptides comprising amino acid sequences sufficiently homologous to or derived  
20 from the amino acid sequence of the TOLL protein, e.g., the amino acid sequence shown in SEQ ID NO:2, which include less amino acids than the full length TOLL proteins, and exhibit at least one activity of a TOLL protein. Typically, biologically active  
35 portions comprise a domain or motif with at least one activity of the TOLL protein. A biologically active portion of a TOLL protein can be a polypeptide which is, for  
40 example, 10, 25, 50, 100, 200 or more amino acids in length. Biologically active  
25 portions of a TOLL protein can be used as targets for developing agents which modulate a TOLL mediated activity.

45 In one embodiment, a biologically active portion of a TOLL protein comprises at least one transmembrane domain, and/or at least one leucine-rich repeat domain and/or  
30 at least one domain containing a  $\beta/\alpha$ -class fold with  $\alpha$ -helices on both faces of a parallel  $\beta$  sheet. It is to be understood that a preferred biologically active portion of a TOLL  
50

5 protein of the present invention may contain at least one transmembrane domain.

Another preferred biologically active portion of a TOLL protein may contain at least one  
10 leucine-rich repeat domain. Another preferred biologically active portion of a TOLL  
protein may contain at least one domain containing a  $\beta/\alpha$ -class fold with  $\alpha$ -helices on  
5 both faces of a parallel  $\beta$  sheet. Moreover, other biologically active portions, in which  
other regions of the protein are deleted, can be prepared by recombinant techniques and  
15 evaluated for one or more of the functional activities of a native TOLL protein.

In a preferred embodiment, the TOLL protein has an amino acid sequence shown  
in SEQ ID NO:2. In other embodiments, the TOLL protein is substantially homologous  
20 to SEQ ID NO:2, and retains the functional activity of the protein of SEQ ID NO:2, yet  
differs in amino acid sequence due to natural allelic variation or mutagenesis, as  
described in detail in subsection I above. Accordingly, in another embodiment, the  
TOLL protein is a protein which comprises an amino acid sequence at least about 50%,  
25 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or more homologous to SEQ  
15 ID NO:2.

To determine the percent identity of two amino acid sequences or of two nucleic  
30 acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps  
can be introduced in one or both of a first and a second amino acid or nucleic acid  
sequence for optimal alignment and non-homologous sequences can be disregarded for  
20 comparison purposes). In a preferred embodiment, the length of a reference sequence  
35 aligned for comparison purposes is at least 30%, preferably at least 40%, more  
preferably at least 50%, even more preferably at least 60%, and even more preferably at  
least 70%, 80%, or 90% of the length of the reference sequence (e.g., when aligning a  
40 second sequence to the TOLL amino acid sequence of SEQ ID NO:2 having 548 amino  
25 acid residues, at least 250, preferably at least 300, more preferably at least 350, even  
more preferably at least 400, and even more preferably at least 450 or 500 amino acid  
residues are aligned). The amino acid residues or nucleotides at corresponding amino  
45 acid positions or nucleotide positions are then compared. When a position in the first  
sequence is occupied by the same amino acid residue or nucleotide as the corresponding  
30 position in the second sequence, then the molecules are identical at that position (as used  
50 herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid

5 "homology"). The percent identity between the two sequences is a function of the  
number of identical positions shared by the sequences, taking into account the number of  
10 gaps, and the length of each gap, which need to be introduced for optimal alignment of  
the two sequences.

5 The comparison of sequences and determination of percent identity between two  
sequences can be accomplished using a mathematical algorithm. In a preferred  
15 embodiment, the percent identity between two amino acid sequences is determined using  
the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has  
been incorporated into the GAP program in the GCG software package (available at  
10 <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap  
weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet  
20 another preferred embodiment, the percent identity between two nucleotide sequences is  
determined using the GAP program in the GCG software package (available at  
25 <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60,  
15 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent  
identity between two amino acid or nucleotide sequences is determined using the  
algorithm of E. Meyers and W. Miller (*CABIOS*, 4:11-17 (1989)) which has been  
30 incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue  
table, a gap length penalty of 12 and a gap penalty of 4.

20 The nucleic acid and protein sequences of the present invention can further be  
35 used as a "query sequence" to perform a search against public databases to, for example,  
identify other family members or related sequences. Such searches can be performed  
using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. (1990) *J.*  
40 *Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the  
25 NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences  
homologous to TOLL nucleic acid molecules of the invention. BLAST protein searches  
can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain  
45 amino acid sequences homologous to TOLL protein molecules of the invention. To  
obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as  
30 described in Altschul et al., (1997) *Nucleic Acids Res.* 25(17):3389-3402. When

utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

The invention also provides TOLL chimeric or fusion proteins. As used herein, a TOLL "chimeric protein" or "fusion protein" comprises a TOLL polypeptide operatively linked to a non-TOLL polypeptide. A "TOLL polypeptide" refers to a polypeptide having an amino acid sequence corresponding to TOLL, whereas a "non-TOLL polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the TOLL protein, e.g., a protein which is different from the TOLL protein and which is derived from the same or a different organism. Within a TOLL fusion protein the TOLL polypeptide can correspond to all or a portion of a TOLL protein. In a preferred embodiment, a TOLL fusion protein comprises at least one biologically active portion of a TOLL protein. In another preferred embodiment, a TOLL fusion protein comprises at least two biologically active portions of a TOLL protein. Within the fusion protein, the term "operatively linked" is intended to indicate that the TOLL polypeptide and the non-TOLL polypeptide are fused in-frame to each other. The non-TOLL polypeptide can be fused to the N-terminus or C-terminus of the TOLL polypeptide.

For example, in one embodiment, the fusion protein is a GST-TOLL fusion protein in which the TOLL sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant TOLL.

In another embodiment, the fusion protein is a TOLL protein containing a heterologous signal sequence at its N-terminus. In certain host cells (e.g., mammalian host cells), expression and/or secretion of TOLL can be increased through use of a heterologous signal sequence.

The TOLL fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject *in vivo*. The TOLL fusion proteins can be used to affect the bioavailability of a TOLL substrate. Use of TOLL fusion proteins may be useful therapeutically for the treatment of disorders caused by, for example, (i) aberrant modification or mutation of a gene encoding a TOLL protein; (ii) mis-regulation of the TOLL gene; and (iii) aberrant post-translational modification of a TOLL protein.

Moreover, the TOLL-fusion proteins of the invention can be used as immunogens to produce anti-TOLL antibodies in a subject, to purify TOLL ligands and in screening assays to identify molecules which inhibit the interaction of TOLL with a TOLL substrate.

Preferably, a TOLL chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST polypeptide). A TOLL-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the TOLL protein.

The present invention also pertains to variants of the TOLL proteins which function as either TOLL agonists (mimetics) or as TOLL antagonists. Variants of the TOLL proteins can be generated by mutagenesis, e.g., discrete point mutation or truncation of a TOLL protein. An agonist of the TOLL proteins can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of a TOLL protein. An antagonist of a TOLL protein can inhibit one or more of the activities of the naturally occurring form of the TOLL protein by, for example, competitively modulating a TOLL-mediated activity of a TOLL protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological

5 activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the TOLL protein.

10 In one embodiment, variants of a TOLL protein which function as either TOLL agonists (mimetics) or as TOLL antagonists can be identified by screening  
5 combinatorial libraries of mutants, e.g., truncation mutants, of a TOLL protein for TOLL protein agonist or antagonist activity. In one embodiment, a variegated library of  
15 TOLL variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of TOLL variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides  
20 into gene sequences such that a degenerate set of potential TOLL sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (e.g., for phage display) containing the set of TOLL sequences therein. There are a variety of methods which can be used to produce libraries of potential TOLL variants  
25 from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes  
30 allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential TOLL sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, e.g., Narang, S.A. (1983) *Tetrahedron* 39:3; Itakura et al. (1984) *Annu. Rev. Biochem.* 53:323; Itakura et al. (1984) *Science* 198:1056; Ike et al. (1983) *Nucleic Acid Res.* 11:477.  
35

In addition, libraries of fragments of a TOLL protein coding sequence can be used to generate a variegated population of TOLL fragments for screening and  
40 subsequent selection of variants of a TOLL protein. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR  
25 fragment of a TOLL coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from  
45 different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression  
30  
50

5 vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of the TOLL protein.

10 Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA  
5 libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of  
15 TOLL proteins. The most widely used techniques, which are amenable to high throughput analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting  
20 library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in  
25 combination with the screening assays to identify TOLL variants (Arkin and Yourvan  
15 (1992) *Proc. Natl. Acad. Sci. USA* 89:7811-7815; Delgrave et al. (1993) *Protein Engineering* 6(3):327-331).

30 In one embodiment, cell based assays can be exploited to analyze a variegated TOLL library. For example, a library of expression vectors can be transfected into a cell line which ordinarily responds to a particular ligand in a TOLL-dependent manner. The  
20 transfected cells are then contacted with the ligand and the effect of expression of the mutant on signaling by the ligand can be detected, e.g., by measuring the activity of a  
35 TOLL-regulated transcription factor. Plasmid DNA can then be recovered from the cells which score for inhibition, or alternatively, potentiation of signaling by the ligand, and the individual clones further characterized.

40 An isolated TOLL protein, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind TOLL using standard techniques for polyclonal and monoclonal antibody preparation. A full-length TOLL protein can be  
45 used or, alternatively, the invention provides antigenic peptide fragments of TOLL for use as immunogens. The antigenic peptide of TOLL comprises at least 8 amino acid  
30 residues of the amino acid sequence shown in SEQ ID NO:2 and encompasses an epitope of TOLL such that an antibody raised against the peptide forms a specific  
50



immune complex with TOLL. Preferably, the antigenic peptide comprises at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at least 30 amino acid residues.

Preferred epitopes encompassed by the antigenic peptide are regions of TOLL that are located on the surface of the protein, e.g., hydrophilic regions, as well as regions with high antigenicity.

A TOLL immunogen typically is used to prepare antibodies by immunizing a suitable subject, (e.g., rabbit, goat, mouse or other mammal) with the immunogen. An appropriate immunogenic preparation can contain, for example, recombinantly expressed TOLL protein or a chemically synthesized TOLL polypeptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic TOLL preparation induces a polyclonal anti-TOLL antibody response.

Accordingly, another aspect of the invention pertains to anti-TOLL antibodies. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site which specifically binds (immunoreacts with) an antigen, such as TOLL. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and F(ab')<sub>2</sub> fragments which can be generated by treating the antibody with an enzyme such as pepsin. The invention provides polyclonal and monoclonal antibodies that bind TOLL. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of TOLL. A monoclonal antibody composition thus typically displays a single binding affinity for a particular TOLL protein with which it immunoreacts.

Polyclonal anti-TOLL antibodies can be prepared as described above by immunizing a suitable subject with a TOLL immunogen. The anti-TOLL antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized TOLL. If

5 desired, the antibody molecules directed against TOLL can be isolated from the mammal  
(e.g., from the blood) and further purified by well known techniques, such as protein A  
10 chromatography to obtain the IgG fraction. At an appropriate time after immunization,  
e.g., when the anti-TOLL antibody titers are highest, antibody-producing cells can be  
5 obtained from the subject and used to prepare monoclonal antibodies by standard  
techniques, such as the hybridoma technique originally described by Kohler and  
15 Milstein (1975) *Nature* 256:495-497) (see also, Brown et al. (1981) *J. Immunol.*  
127:539-46; Brown et al. (1980) *J. Biol. Chem.* 255:4980-83; Yeh et al. (1976) *Proc.*  
*Natl. Acad. Sci. USA* 76:2927-31; and Yeh et al. (1982) *Int. J. Cancer* 29:269-75), the  
20 more recent human B cell hybridoma technique (Kozbor et al. (1983) *Immunol Today*  
4:72), the EBV-hybridoma technique (Cole et al. (1985), *Monoclonal Antibodies and*  
*Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology  
for producing monoclonal antibody hybridomas is well known (see generally R. H.  
25 Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum  
15 Publishing Corp., New York, New York (1980); E. A. Lerner (1981) *Yale J. Biol. Med.*,  
54:387-402; M. L. Gefter et al. (1977) *Somatic Cell Genet.* 3:231-36). Briefly, an  
immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes)  
30 from a mammal immunized with a TOLL immunogen as described above, and the  
culture supernatants of the resulting hybridoma cells are screened to identify a  
20 hybridoma producing a monoclonal antibody that binds TOLL.

35 Any of the many well known protocols used for fusing lymphocytes and  
immortalized cell lines can be applied for the purpose of generating an anti-TOLL  
monoclonal antibody (see, e.g., G. Galfre et al. (1977) *Nature* 266:55052; Gefter et al.  
40 *Somatic Cell Genet.*, cited *supra*; Lerner, *Yale J. Biol. Med.*, cited *supra*; Kenneth,  
25 *Monoclonal Antibodies*, cited *supra*). Moreover, the ordinarily skilled worker will  
appreciate that there are many variations of such methods which also would be useful.  
Typically, the immortal cell line (e.g., a myeloma cell line) is derived from the same  
45 mammalian species as the lymphocytes. For example, murine hybridomas can be made  
by fusing lymphocytes from a mouse immunized with an immunogenic preparation of  
30 the present invention with an immortalized mouse cell line. Preferred immortal cell  
50 lines are mouse myeloma cell lines that are sensitive to culture medium containing

5 hypoxanthine, aminopterin and thymidine ("HAT medium"). Any of a number of  
myeloma cell lines can be used as a fusion partner according to standard techniques,  
e.g., the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/O-Ag14 myeloma lines. These  
10 myeloma lines are available from ATCC. Typically, HAT-sensitive mouse myeloma  
5 cells are fused to mouse splenocytes using polyethylene glycol ("PEG"). Hybridoma  
cells resulting from the fusion are then selected using HAT medium, which kills unfused  
and unproductively fused myeloma cells (unfused splenocytes die after several days  
15 because they are not transformed). Hybridoma cells producing a monoclonal antibody  
of the invention are detected by screening the hybridoma culture supernatants for  
20 antibodies that bind TOLL, e.g., using a standard ELISA assay.

Alternative to preparing monoclonal antibody-secreting hybridomas, a  
monoclonal anti-TOLL antibody can be identified and isolated by screening a  
recombinant combinatorial immunoglobulin library (e.g., an antibody phage display  
25 library) with TOLL to thereby isolate immunoglobulin library members that bind TOLL.  
15 Kits for generating and screening phage display libraries are commercially available  
(e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and  
the Stratagene *SurfZAP™ Phage Display Kit*, Catalog No. 240612). Additionally,  
30 examples of methods and reagents particularly amenable for use in generating and  
screening antibody display library can be found in, for example, Ladner et al. U.S.  
20 Patent No. 5,223,409; Kang et al. PCT International Publication No. WO 92/18619;  
35 Dower et al. PCT International Publication No. WO 91/17271; Winter et al. PCT  
International Publication WO 92/20791; Markland et al. PCT International Publication  
No. WO 92/15679; Breitling et al. PCT International Publication WO 93/01288;  
40 McCafferty et al. PCT International Publication No. WO 92/01047; Garrard et al. PCT  
25 International Publication No. WO 92/09690; Ladner et al. PCT International Publication  
No. WO 90/02809; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992)  
*Hum. Antibod. Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-1281;  
45 Griffiths et al. (1993) *EMBO J* 12:725-734; Hawkins et al. (1992) *J. Mol. Biol.* 226:889-  
896; Clarkson et al. (1991) *Nature* 352:624-628; Gram et al. (1992) *Proc. Natl. Acad.*  
30 *Sci. USA* 89:3576-3580; Garrad et al. (1991) *Bio/Technology* 9:1373-1377;

5 Hoogenboom et al. (1991) *Nuc. Acid Res.* 19:4133-4137; Barbas et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:7978-7982; and McCafferty et al. *Nature* (1990) 348:552-554.

10 Additionally, recombinant anti-TOLL antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in Robinson et al. International Application No. PCT/US86/02269; Akira, et al. European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison et al. European Patent Application 173,494; Neuberger et al. PCT International Publication No. WO 86/01533; Cabilly et al. U.S. Patent No. 4,816,567; Cabilly et al. European Patent Application 125,023; Better et al. (1988) *Science* 240:1041-1043; Liu et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:3439-3443; Liu et al. (1987) *J. Immunol.* 139:3521-3526; Sun et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:214-218; Nishimura et al. (1987) *Canc. Res.* 47:999-1005; Wood et al. (1985) *Nature* 314:446-449; and Shaw et al. (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison, S. L. (1985) *Science* 229:1202-1207; Oi et al. (1986) *BioTechniques* 4:214; Winter U.S. Patent 5,225,539; Jones et al. (1986) *Nature* 321:552-525; Verhoeyan et al. (1988) *Science* 239:1534; and Beidler et al. (1988) *J. Immunol.* 141:4053-4060.

20 An anti-TOLL antibody (e.g., monoclonal antibody) can be used to isolate TOLL proteins by standard techniques, such as affinity chromatography or immunoprecipitation. An anti-TOLL antibody can facilitate the purification of natural TOLL proteins from cells and of recombinantly produced TOLL proteins expressed in host cells. Moreover, an anti-TOLL antibody can be used to detect TOLL protein (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the TOLL protein. Anti-TOLL antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, e.g., to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive

materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$  or  $^3\text{H}$ .

### III. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding a TOLL protein (or a portion thereof). As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

5 The recombinant expression vectors of the invention comprise a nucleic acid of  
the invention in a form suitable for expression of the nucleic acid in a host cell, which  
means that the recombinant expression vectors include one or more regulatory  
10 sequences, selected on the basis of the host cells to be used for expression, which is  
5 operatively linked to the nucleic acid sequence to be expressed. Within a recombinant  
expression vector, "operably linked" is intended to mean that the nucleotide sequence of  
interest is linked to the regulatory sequence(s) in a manner which allows for expression  
15 of the nucleotide sequence (e.g., in an *in vitro* transcription/translation system or in a  
host cell when the vector is introduced into the host cell). The term "regulatory  
20 sequence" is intended to include promoters, enhancers and other expression control  
elements (e.g., polyadenylation signals). Such regulatory sequences are described, for  
example, in Goeddel: *Gene Expression Technology: Methods in Enzymology* 185,  
Academic Press, San Diego, CA (1990). Regulatory sequences include those which  
25 direct constitutive expression of a nucleotide sequence in many types of host cells and  
15 those which direct expression of the nucleotide sequence only in certain host cells (e.g.,  
tissue-specific regulatory sequences). It will be appreciated by those skilled in the art  
that the design of the expression vector can depend on such factors as the choice of the  
30 host cell to be transformed, the level of expression of protein desired, and the like. The  
expression vectors of the invention can be introduced into host cells to thereby produce  
20 proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as  
described herein (e.g., TOLL proteins, mutant forms of TOLL proteins, fusion proteins,  
35 and the like).

The recombinant expression vectors of the invention can be designed for  
40 expression of TOLL proteins in prokaryotic or eukaryotic cells. For example, TOLL  
25 proteins can be expressed in bacterial cells such as *E. coli*, insect cells (using  
baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are  
discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology*  
45 185, Academic Press, San Diego, CA (1990). Alternatively, the recombinant expression  
vector can be transcribed and translated *in vitro*, for example using T7 promoter  
30 regulatory sequences and T7 polymerase.

5 Expression of proteins in prokaryotes is most often carried out in *E. coli* with  
vectors containing constitutive or inducible promoters directing the expression of either  
fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein  
10 encoded therein, usually to the amino terminus of the recombinant protein. Such fusion  
vectors typically serve three purposes: 1) to increase expression of recombinant protein;  
2) to increase the solubility of the recombinant protein; and 3) to aid in the purification  
15 of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion  
expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion  
moiety and the recombinant protein to enable separation of the recombinant protein from  
20 the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and  
their cognate recognition sequences, include Factor Xa, thrombin and enterokinase.  
Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B.  
and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA)  
25 and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST),  
maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Purified fusion proteins can be utilized in TOLL activity assays, (e.g., direct  
assays or competitive assays described in detail below), or to generate antibodies  
30 specific for TOLL proteins, for example. In a preferred embodiment, a TOLL fusion  
protein expressed in a retroviral expression vector of the present invention can be  
utilized to infect bone marrow cells which are subsequently transplanted into irradiated  
20 recipients. The pathology of the subject recipient is then examined after sufficient time  
has passed (e.g., six (6) weeks).

Examples of suitable inducible non-fusion *E. coli* expression vectors include  
pTrc (Amann et al., (1988) *Gene* 69:301-315) and pET 11d (Studier et al., *Gene*  
40 *Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego,  
California (1990) 60-89). Target gene expression from the pTrc vector relies on host  
RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene  
45 expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion  
promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral  
polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident  
30

5 prophage harboring a T7 *gn1* gene under the transcriptional control of the lacUV 5 promoter.

10 One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in*  
15 *Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada et al., (1992) *Nucleic Acids Res.* 20:2111-2118).  
20 Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the TOLL expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, et  
25 al., (1987) *EMBO J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz et al., (1987) *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, CA), and picZ (Invitrogen Corp, San Diego, CA).

30 Alternatively, TOLL proteins can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith et al. (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

35 In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman et al. (1987) *EMBO J.* 6:187-195). When used in mammalian cells, the  
40 expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both  
45 prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY,  
50 1989.



5 In another embodiment, the recombinant mammalian expression vector is  
capable of directing expression of the nucleic acid preferentially in a particular cell type  
(e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-  
10 specific regulatory elements are known in the art. Non-limiting examples of suitable  
5 tissue-specific promoters include the albumin promoter (liver-specific; Pinkert et al.  
(1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988)  
*Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winkler and  
15 Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji et al. (1983) *Cell*  
33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters  
10 (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *Proc. Natl. Acad. Sci. USA*  
86:5473-5477), pancreas-specific promoters (Edlund et al. (1985) *Science* 230:912-916),  
20 and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No.  
4,873,316 and European Application Publication No. 264,166). Developmentally-  
25 regulated promoters are also encompassed, for example the murine hox promoters  
15 (Kessel and Gruss (1990) *Science* 249:374-379) and the  $\alpha$ -fetoprotein promoter  
(Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

30 The invention further provides a recombinant expression vector comprising a  
DNA molecule of the invention cloned into the expression vector in an antisense-  
orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in  
20 a manner which allows for expression (by transcription of the DNA molecule) of an  
RNA molecule which is antisense to TOLL mRNA. Regulatory sequences operatively  
35 linked to a nucleic acid cloned in the antisense orientation can be chosen which direct  
the continuous expression of the antisense RNA molecule in a variety of cell types, for  
instance viral promoters and/or enhancers, or regulatory sequences can be chosen which  
40 25 direct constitutive, tissue specific or cell type specific expression of antisense RNA. The  
antisense expression vector can be in the form of a recombinant plasmid, phagemid or  
attenuated virus in which antisense nucleic acids are produced under the control of a  
45 high efficiency regulatory region, the activity of which can be determined by the cell  
type into which the vector is introduced. For a discussion of the regulation of gene  
30 expression using antisense genes see Weintraub, H. et al., Antisense RNA as a molecular  
50 tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a TOLL nucleic acid molecule of the invention is introduced, e.g., a TOLL nucleic acid molecule within a recombinant expression vector or a TOLL nucleic acid molecule containing sequences which allow it to homologously recombine into a specific site of the host-cell's genome.

The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, a TOLL protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, et al. (*Molecular Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding a TOLL protein or can be

5 introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

10 A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (i.e., express) a TOLL protein. Accordingly, the invention further provides methods for producing a TOLL protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of the invention (into which a recombinant expression vector encoding a TOLL protein has been introduced) in a suitable medium such that a TOLL protein is produced. In another 15 embodiment, the method further comprises isolating a TOLL protein from the medium or the host cell.

The host cells of the invention can also be used to produce non-human transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which TOLL-coding sequences have been 25 introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous TOLL sequences have been introduced into their genome or homologous recombinant animals in which endogenous TOLL sequences have been altered. Such animals are useful for studying the function and/or activity of a TOLL and for identifying and/or evaluating modulators of TOLL activity. As used herein, a 30 "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, and the like. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops 35 and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous TOLL gene has been altered by homologous recombination between the endogenous gene and an exogenous 40 DNA molecule introduced into a cell of the animal, e.g., an embryonic cell of the animal, prior to development of the animal.

5 A transgenic animal of the invention can be created by introducing a TOLL-  
encoding nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by  
microinjection, retroviral infection, and allowing the oocyte to develop in a  
10 pseudopregnant female foster animal. The TOLL cDNA sequence of SEQ ID NO:1 can  
5 be introduced as a transgene into the genome of a non-human animal. Alternatively, a  
nonhuman homologue of a human TOLL gene, such as a mouse or rat TOLL gene, or  
the sequence of SEQ ID NO:4 can be used as a transgene. Alternatively, a TOLL gene  
15 homologue, such as another TOLL family member, can be isolated based on  
hybridization to the TOLL cDNA sequences of SEQ ID NO:1, 3, or 4, or the DNA insert  
10 of the plasmid deposited with ATCC as Accession Number \_\_\_\_\_ (described further in  
subsection I above) and used as a transgene. Intronic sequences and polyadenylation  
signals can also be included in the transgene to increase the efficiency of expression of  
the transgene. A tissue-specific regulatory sequence(s) can be operably linked to a  
25 TOLL transgene to direct expression of a TOLL protein to particular cells. Methods for  
15 generating transgenic animals via embryo manipulation and microinjection, particularly  
animals such as mice, have become conventional in the art and are described, for  
example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder et al., U.S. Patent  
30 No. 4,873,191 by Wagner et al. and in Hogan, B., *Manipulating the Mouse Embryo*,  
(Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar  
20 methods are used for production of other transgenic animals. A transgenic founder  
animal can be identified based upon the presence of a TOLL transgene in its genome  
and/or expression of TOLL mRNA in tissues or cells of the animals. A transgenic  
35 founder animal can then be used to breed additional animals carrying the transgene.  
Moreover, transgenic animals carrying a transgene encoding a TOLL protein can further  
40 be bred to other transgenic animals carrying other transgenes.

To create a homologous recombinant animal, a vector is prepared which contains  
at least a portion of a TOLL gene into which a deletion, addition or substitution has been  
45 introduced to thereby alter, e.g., functionally disrupt, the TOLL gene. The TOLL gene  
can be a human gene (e.g., the cDNA of SEQ ID NO:1), but more preferably, is a non-  
30 human homologue of a human TOLL gene (e.g., a cDNA isolated by stringent  
50 hybridization with the nucleotide sequence of SEQ ID NO:1, or the sequence of SEQ ID

5 NO: 4). For example, a mouse TOLL gene can be used to construct a homologous  
recombination nucleic acid molecule, e.g., a vector, suitable for altering an endogenous  
10 TOLL gene in the mouse genome. In a preferred embodiment, the homologous  
recombination nucleic acid molecule is designed such that, upon homologous  
5 recombination, the endogenous TOLL gene is functionally disrupted (i.e., no longer  
encodes a functional protein; also referred to as a "knock out" vector). Alternatively, the  
homologous recombination nucleic acid molecule can be designed such that, upon  
15 homologous recombination, the endogenous TOLL gene is mutated or otherwise altered  
but still encodes functional protein (e.g., the upstream regulatory region can be altered to  
20 thereby alter the expression of the endogenous TOLL protein). In the homologous  
recombination nucleic acid molecule, the altered portion of the TOLL gene is flanked at  
its 5' and 3' ends by additional nucleic acid sequence of the TOLL gene to allow for  
homologous recombination to occur between the exogenous TOLL gene carried by the  
25 homologous recombination nucleic acid molecule and an endogenous TOLL gene in a  
cell, e.g., an embryonic stem cell. The additional flanking TOLL nucleic acid sequence  
is of sufficient length for successful homologous recombination with the endogenous  
30 gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are  
included in the homologous recombination nucleic acid molecule (see, e.g., Thomas,  
K.R. and Capecchi, M. R. (1987) *Cell* 51:503 for a description of homologous  
20 recombination vectors). The homologous recombination nucleic acid molecule is  
introduced into a cell, e.g., an embryonic stem cell line (e.g., by electroporation) and  
35 cells in which the introduced TOLL gene has homologously recombined with the  
endogenous TOLL gene are selected (see e.g., Li, E. et al. (1992) *Cell* 69:915). The  
selected cells can then be injected into a blastocyst of an animal (e.g., a mouse) to form  
40 aggregation chimeras (see e.g., Bradley, A. in *Teratocarcinomas and Embryonic Stem  
Cells: A Practical Approach*, E.J. Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A  
chimeric embryo can then be implanted into a suitable pseudopregnant female foster  
45 animal and the embryo brought to term. Progeny harboring the homologously  
recombined DNA in their germ cells can be used to breed animals in which all cells of  
30 the animal contain the homologously recombined DNA by germline transmission of the  
transgene. Methods for constructing homologous recombination nucleic acid molecules,

5 e.g., vectors, or homologous recombinant animals are described further in Bradley, A.  
(1991) *Current Opinion in Biotechnology* 2:823-829 and in PCT International  
Publication Nos.: WO 90/11354 by Le Mouellec et al.; WO 91/01140 by Smithies et  
10 al.; WO 92/0968 by Zijlstra et al.; and WO 93/04169 by Berns et al.

5 In another embodiment, transgenic non-humans animals can be produced which  
contain selected systems which allow for regulated expression of the transgene. One  
example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For  
15 a description of the *cre/loxP* recombinase system, see, e.g., Iakso et al. (1992) *Proc.*  
*Natl. Acad. Sci. USA* 89:6232-6236. Another example of a recombinase system is the  
20 FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman et al. (1991) *Science*  
251:1351-1355. If a *cre/loxP* recombinase system is used to regulate expression of the  
transgene, animals containing transgenes encoding both the *Cre* recombinase and a  
selected protein are required. Such animals can be provided through the construction of  
25 "double" transgenic animals, e.g., by mating two transgenic animals, one containing a  
15 transgene encoding a selected protein and the other containing a transgene encoding a  
recombinase.

30 Clones of the non-human transgenic animals described herein can also be  
produced according to the methods described in Wilmut, I. et al. (1997) *Nature* 385:810-  
813 and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief,  
20 a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit  
35 the growth cycle and enter G<sub>0</sub> phase. The quiescent cell can then be fused, e.g., through  
the use of electrical pulses, to an enucleated oocyte from an animal of the same species  
from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such  
40 that it develops to morula or blastocyte and then transferred to pseudopregnant female  
25 foster animal. The offspring borne of this female foster animal will be a clone of the  
animal from which the cell, e.g., the somatic cell, is isolated.

#### 45 IV. Pharmaceutical Compositions

The TOLL nucleic acid molecules, fragments of TOLL proteins, and anti-TOLL  
30 antibodies (also referred to herein as "active compounds") of the invention can be  
50 incorporated into pharmaceutical compositions suitable for administration. Such

5 compositions typically comprise the nucleic acid molecule, protein, or antibody and a  
pharmaceutically acceptable carrier. As used herein the language "pharmaceutically  
10 acceptable carrier" is intended to include any and all solvents, dispersion media,  
coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents,  
5 and the like, compatible with pharmaceutical administration. The use of such media and  
agents for pharmaceutically active substances is well known in the art. Except insofar as  
15 any conventional media or agent is incompatible with the active compound, use thereof  
in the compositions is contemplated. Supplementary active compounds can also be  
incorporated into the compositions.

10 A pharmaceutical composition of the invention is formulated to be compatible  
with its intended route of administration. Examples of routes of administration include  
20 parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation),  
transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions  
25 used for parenteral, intradermal, or subcutaneous application can include the following  
15 components: a sterile diluent such as water for injection, saline solution, fixed oils,  
polyethylene glycols, glycerine, propylene glycol or other synthetic solvents;  
antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as  
30 ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic  
acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of  
20 tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases,  
35 such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be  
enclosed in ampoules, disposable syringes or multiple dose vials made of glass or  
plastic.

40 Pharmaceutical compositions suitable for injectable use include sterile aqueous  
25 solutions (where water soluble) or dispersions and sterile powders for the  
extemporaneous preparation of sterile injectable solutions or dispersion. For  
intravenous administration, suitable carriers include physiological saline, bacteriostatic  
45 water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In  
all cases, the composition must be sterile and should be fluid to the extent that easy  
30 syringability exists. It must be stable under the conditions of manufacture and storage  
and must be preserved against the contaminating action of microorganisms such as

5 bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for  
example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid  
polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity  
10 can be maintained, for example, by the use of a coating such as lecithin, by the  
maintenance of the required particle size in the case of dispersion and by the use of  
surfactants. Prevention of the action of microorganisms can be achieved by various  
antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol,  
15 ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include  
isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium  
chloride in the composition. Prolonged absorption of the injectable compositions can be  
20 brought about by including in the composition an agent which delays absorption, for  
example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active  
25 compound (e.g., a fragment of a TOLL protein or an anti-TOLL antibody) in the  
required amount in an appropriate solvent with one or a combination of ingredients  
enumerated above, as required, followed by filtered sterilization. Generally, dispersions  
are prepared by incorporating the active compound into a sterile vehicle which contains  
30 a basic dispersion medium and the required other ingredients from those enumerated  
above. In the case of sterile powders for the preparation of sterile injectable solutions,  
the preferred methods of preparation are vacuum drying and freeze-drying which yields  
20 a powder of the active ingredient plus any additional desired ingredient from a  
previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They  
can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral  
40 therapeutic administration, the active compound can be incorporated with excipients and  
used in the form of tablets, troches, or capsules. oral compositions can also be prepared  
using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is  
applied orally and swished and expectorated or swallowed. Pharmaceutically  
45 compatible binding agents, and/or adjuvant materials can be included as part of the  
composition. The tablets, pills, capsules, troches and the like can contain any of the  
50 following ingredients, or compounds of a similar nature: a binder such as



5 microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or  
lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant  
10 such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a  
sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint,  
5 methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an  
aerosol spray from pressured container or dispenser which contains a suitable propellant,  
15 e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For  
10 transmucosal or transdermal administration, penetrants appropriate to the barrier to be  
permeated are used in the formulation. Such penetrants are generally known in the art,  
20 and include, for example, for transmucosal administration, detergents, bile salts, and  
fusidic acid derivatives. Transmucosal administration can be accomplished through the  
use of nasal sprays or suppositories. For transdermal administration, the active  
25 compounds are formulated into ointments, salves, gels, or creams as generally known in  
the art.

The compounds can also be prepared in the form of suppositories (e.g., with  
30 conventional suppository bases such as cocoa butter and other glycerides) or retention  
enemas for rectal delivery.

20 In one embodiment, the active compounds are prepared with carriers that will  
protect the compound against rapid elimination from the body, such as a controlled  
35 release formulation, including implants and microencapsulated delivery systems.  
Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate,  
polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid.  
40 Methods for preparation of such formulations will be apparent to those skilled in the art.  
The materials can also be obtained commercially from Alza Corporation and Nova  
Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected  
45 cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically  
acceptable carriers. These can be prepared according to methods known to those skilled  
30 in the art, for example, as described in U.S. Patent No. 4,522,811.

5 It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound  
10 calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

15 Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio  
20 LD50/ED50. Compounds which exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

25 The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially  
30 from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

5 The nucleic acid molecules of the invention can be inserted into vectors and used  
as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for  
example, intravenous injection, local administration (see U.S. Patent 5,328,470) or by  
10 stereotactic injection (see e.g., Chen et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3054-  
5 3057). The pharmaceutical preparation of the gene therapy vector can include the gene  
therapy vector in an acceptable diluent, or can comprise a slow release matrix in which  
the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery  
15 vector can be produced intact from recombinant cells, e.g., retroviral vectors, the  
pharmaceutical preparation can include one or more cells which produce the gene  
20 delivery system.

The pharmaceutical compositions can be included in a container, pack, or  
dispenser together with instructions for administration.

#### 25 V. Uses and Methods of the Invention

15 The nucleic acid molecules, proteins, protein homologues, and antibodies  
described herein can be used in one or more of the following methods: a) screening  
assays; b) predictive medicine (e.g., diagnostic assays, prognostic assays, monitoring  
30 clinical trials, and pharmacogenetics); and c) methods of treatment (e.g., therapeutic and  
prophylactic). As described herein, a TOLL protein of the invention has one or more of  
20 the following activities: (1) it interacts with a non-TOLL protein molecule, e.g., a  
TOLL ligand; (2) it activates a TOLL-dependent signal transduction pathway; and (3) it  
35 modulates signaling mechanisms involved in, for example, an immune response, and,  
thus, can be used to, for example, (1) modulate the interaction with a non-TOLL protein  
molecule; (2) to activate a TOLL-dependent signal transduction pathway; and (3) to  
40 25 modulate signalling mechanisms involved in, for example, an immune response.

The isolated nucleic acid molecules of the invention can be used, for example, to  
express TOLL protein (e.g., via a recombinant expression vector in a host cell in gene  
45 therapy applications), to detect TOLL mRNA (e.g., in a biological sample) or a genetic  
alteration in a TOLL gene, and to modulate TOLL activity, as described further below.

30 The TOLL proteins can be used to treat disorders characterized by insufficient or  
excessive production of a TOLL substrate or production of TOLL inhibitors. In  
50

5 addition, the TOLL proteins can be used to screen for naturally occurring TOLL  
substrates, to screen for drugs or compounds which modulate TOLL activity, as well as  
10 to treat disorders characterized by insufficient or excessive production of TOLL protein  
or production of TOLL protein forms which have decreased, aberrant or unwanted  
5 activity compared to TOLL wild type protein. Moreover, the anti-TOLL antibodies of  
the invention can be used to detect and isolate TOLL proteins, regulate the  
15 bioavailability of TOLL proteins, and modulate TOLL activity.

A. Screening Assays:

10 The invention provides a method (also referred to herein as a "screening assay")  
for identifying modulators, i.e., candidate or test compounds or agents (e.g., peptides,  
20 peptidomimetics, small molecules or other drugs) which bind to TOLL proteins, have a  
stimulatory or inhibitory effect on, for example, TOLL expression or TOLL activity, or  
25 have a stimulatory or inhibitory effect on, for example, the expression or activity of  
15 TOLL substrate.

In one embodiment, the invention provides assays for screening candidate or test  
30 compounds which are substrates of a TOLL protein or polypeptide or biologically active  
portion thereof. In another embodiment, the invention provides assays for screening  
candidate or test compounds which bind to or modulate the activity of a TOLL protein  
20 or polypeptide or biologically active portion thereof. The test compounds of the present  
invention can be obtained using any of the numerous approaches in combinatorial  
35 library methods known in the art, including: biological libraries; spatially addressable  
parallel solid phase or solution phase libraries; synthetic library methods requiring  
deconvolution; the 'one-bead one-compound' library method; and synthetic library  
40 25 methods using affinity chromatography selection. The biological library approach is  
limited to peptide libraries, while the other four approaches are applicable to peptide,  
non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997)  
45 *Anticancer Drug Des.* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the  
30 art, for example in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.*  
50 (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med.*

5 Chem. 37:2678; Cho et al. (1993) *Science* 261:1303; Carrell et al. (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carrell et al. (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop et al. (1994) *J. Med. Chem.* 37:1233.

10 Libraries of compounds may be presented in solution (e.g., Houghten (1992) *Biotechniques* 13:412-421); or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull et al. (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla et al. (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*).

20 In one embodiment, an assay is a cell-based assay in which a cell which expresses a TOLL protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate TOLL activity is determined. Determining the ability of the test compound to modulate TOLL activity  
25 can be accomplished by monitoring, for example, the activity of a TOLL-regulated transcription factor. The cell, for example, can be of mammalian origin.

30 The ability of the test compound to modulate TOLL binding to a substrate or to bind to TOLL can also be determined. Determining the ability of the test compound to modulate TOLL binding to a substrate can be accomplished, for example, by coupling  
35 the TOLL substrate with a radioisotope or enzymatic label such that binding of the TOLL substrate to TOLL can be determined by detecting the labeled TOLL substrate in a complex. Determining the ability of the test compound to bind TOLL can be accomplished, for example, by coupling the compound with a radioisotope or enzymatic  
40 label such that binding of the compound to TOLL can be determined by detecting the labeled TOLL compound in a complex. For example, compounds (e.g., TOLL substrates) can be labeled with  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , or  $^3\text{H}$ , either directly or indirectly, and  
25 the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label  
30 detected by determination of conversion of an appropriate substrate to product.

5 It is also within the scope of this invention to determine the ability of a compound (e.g., a TOLL substrate) to interact with TOLL without the labeling of any of the interactants. For example, a microphysiometer can be used to detect the interaction of a compound with TOLL without the labeling of either the compound or the TOLL.

10  
5 McConnell, H. M. et al. (1992) *Science* 257:1906-1912. As used herein, a "microphysiometer" (e.g., Cytosensor) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a compound and TOLL.

15  
10 In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a TOLL target molecule (e.g., a TOLL substrate) with a test compound and determining the ability of the test compound to modulate (e.g. stimulate or inhibit) the activity of the TOLL target molecule. Determining the ability of the test compound to modulate the activity of a TOLL target molecule can be accomplished, for example, by determining the ability of the TOLL protein to bind to or interact with the TOLL target molecule.

20  
25  
30 Determining the ability of the TOLL protein or a biologically active fragment thereof, to bind to or interact with a TOLL target molecule can be accomplished by one of the methods described above for determining direct binding. In a preferred embodiment, determining the ability of the TOLL protein to bind to or interact with a TOLL target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (e.g., intracellular  $\text{Ca}^{2+}$ , diacylglycerol,  $\text{IP}_3$ , and the like), detecting catalytic/enzymatic activity of the target an appropriate substrate, detecting the induction of a reporter gene (comprising a target-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a target-regulated cellular response.

35  
40  
45  
50 In yet another embodiment, an assay of the present invention is a cell-free assay in which a TOLL protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to the TOLL protein or biologically active portion thereof is determined. Preferred biologically active portions

5 of the TOLL proteins to be used in assays of the present invention include fragments  
which participate in interactions with non-TOLL molecules, e.g., fragments with high  
10 surface probability scores. Binding of the test compound to the TOLL protein can be  
determined either directly or indirectly as described above. In a preferred embodiment,  
5 the assay includes contacting the TOLL protein or biologically active portion thereof  
with a known compound which binds TOLL to form an assay mixture, contacting the  
15 assay mixture with a test compound, and determining the ability of the test compound to  
interact with a TOLL protein, wherein determining the ability of the test compound to  
interact with a TOLL protein comprises determining the ability of the test compound to  
10 preferentially bind to TOLL or biologically active portion thereof as compared to the  
20 known compound.

In another embodiment, the assay is a cell-free assay in which a TOLL protein or  
biologically active portion thereof is contacted with a test compound and the ability of  
25 the test compound to modulate (e.g., stimulate or inhibit) the activity of the TOLL  
protein or biologically active portion thereof is determined. Determining the ability of  
15 the test compound to modulate the activity of a TOLL protein can be accomplished, for  
example, by determining the ability of the TOLL protein to bind to a TOLL target  
30 molecule by one of the methods described above for determining direct binding.  
Determining the ability of the TOLL protein to bind to a TOLL target molecule can also  
20 be accomplished using a technology such as real-time Biomolecular Interaction Analysis  
(BIA). Sjölander, S. and Urbaniczky, C. (1991) *Anal. Chem.* 63:2338-2345 and Szabo  
35 et al. (1995) *Curr. Opin. Struct. Biol.* 5:699-705. As used herein, "BIA" is a technology  
for studying biospecific interactions in real time, without labeling any of the interactants  
(e.g., BIAcore). Changes in the optical phenomenon of surface plasmon resonance  
40 (SPR) can be used as an indication of real-time reactions between biological molecules.  
25

In an alternative embodiment, determining the ability of the test compound to  
modulate the activity of a TOLL protein can be accomplished by determining the ability  
45 of the TOLL protein to further modulate the activity of a downstream effector of a  
TOLL target molecule. For example, the activity of the effector molecule on an  
30 appropriate target can be determined or the binding of the effector to an appropriate  
target can be determined as previously described.  
50

5 In yet another embodiment, the cell-free assay involves contacting a TOLL protein or biologically active portion thereof with a known compound which binds the TOLL protein to form an assay mixture, contacting the assay mixture with a test  
10 compound, and determining the ability of the test compound to interact with the TOLL protein, wherein determining the ability of the test compound to interact with the TOLL protein comprises determining the ability of the TOLL protein to preferentially bind to or modulate the activity of a TOLL target molecule.

15 The cell-free assays of the present invention are amenable to use of both soluble and/or membrane-bound forms of isolated proteins (e.g., TOLL proteins or biologically active portions thereof). In the case of cell-free assays in which a membrane-bound  
20 form of an isolated protein is used it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the isolated protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide,  
25 decanoyl-N-methylglucamide, Triton® X-100, Triton® X-114, Thesit®, Isotridecypoly(ethylene glycol ether)<sub>n</sub>, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl=N,N-dimethyl-3-ammonio-1-propane sulfonate.

20 In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either TOLL or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test  
35 compound to a TOLL protein, or interaction of a TOLL protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be  
40 provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase/ TOLL fusion proteins or glutathione-S-transferase/target fusion proteins can be adsorbed onto glutathione sepharose beads  
30 (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are



5 then combined with the test compound or the test compound and either the non-adsorbed target protein or TOLL protein. and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following  
10 incubation, the beads or microtitre plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads. complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of TOLL binding or activity determined  
15 using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the  
10 screening assays of the invention. For example, either a TOLL protein or a TOLL target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated TOLL protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques known in the art (e.g., biotinylation kit, Pierce  
20 Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with TOLL protein or target molecules but which do not interfere with binding of the TOLL protein to its target molecule can be derivatized to the wells of the plate, and unbound target or TOLL  
25 protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the TOLL protein  
30 or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the TOLL protein or target molecule.

In another embodiment, modulators of TOLL expression are identified in a  
40 method wherein a cell is contacted with a candidate compound and the expression of TOLL mRNA or protein in the cell is determined. The level of expression of TOLL mRNA or protein in the presence of the candidate compound is compared to the level of expression of TOLL mRNA or protein in the absence of the candidate compound. The  
45 candidate compound can then be identified as a modulator of TOLL expression based on this comparison. For example, when expression of TOLL mRNA or protein is greater  
50 (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of TOLL mRNA or

5 protein expression. Alternatively, when expression of TOLL mRNA or protein is less  
(statistically significantly less) in the presence of the candidate compound than in its  
10 absence, the candidate compound is identified as an inhibitor of TOLL mRNA or protein  
expression. The level of TOLL mRNA or protein expression in the cells can be  
5 determined by methods described herein for detecting TOLL mRNA or protein.

In yet another aspect of the invention, the TOLL proteins can be used as "bait  
15 proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No.  
5,283,317; Zervos et al. (1993) *Cell* 72:223-232; Madura et al. (1993) *J. Biol. Chem.*  
268:12046-12054; Bartel et al. (1993) *Biotechniques* 14:920-924; Iwabuchi et al. (1993)  
20 *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind  
to or interact with TOLL ("TOLL-binding proteins" or "TOLL-bp") and are involved in  
TOLL activity. Such TOLL-binding proteins are also likely to be involved in the  
propagation of signals by the TOLL proteins or TOLL targets as, for example,  
25 downstream elements of a TOLL-mediated signaling pathway. Alternatively, such  
15 TOLL-binding proteins are likely to be TOLL inhibitors.

The two-hybrid system is based on the modular nature of most transcription  
factors, which consist of separable DNA-binding and activation domains. Briefly, the  
30 assay utilizes two different DNA constructs. In one construct, the gene that codes for a  
TOLL protein is fused to a gene encoding the DNA binding domain of a known  
20 transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a  
library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is  
35 fused to a gene that codes for the activation domain of the known transcription factor. If  
the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a TOLL-  
dependent complex, the DNA-binding and activation domains of the transcription factor  
40 are brought into close proximity. This proximity allows transcription of a reporter gene  
25 (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the  
transcription factor. Expression of the reporter gene can be detected and cell colonies  
45 containing the functional transcription factor can be isolated and used to obtain the  
cloned gene which encodes the protein which interacts with the TOLL protein.

5 In another aspect, the invention pertains to a combination of two or more of the assays described herein. For example, a modulating agent can be identified using a cell-based or a cell free assay, and the ability of the agent to modulate the activity of a TOLL  
10 protein can be confirmed *in vivo*, e.g., in an animal such as an animal model for a TOLL associated disorder.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an  
15 agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a TOLL modulating agent, an antisense TOLL nucleic acid molecule, a TOLL-specific antibody, or a TOLL-binding partner) can be  
20 used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore,  
25 this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

#### 30 B. Detection Assays

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as  
20 polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with  
35 genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

##### 40 25 1. Chromosome Mapping

Once the sequence (or a portion of the sequence) of a gene has been isolated, this  
45 sequence can be used to map the location of the gene on a chromosome. This process is called chromosome mapping. Accordingly, portions or fragments of the TOLL  
30 nucleotide sequences, described herein, can be used to map the location of the TOLL

genes on a chromosome. The mapping of the TOLL sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Briefly, TOLL genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the TOLL nucleotide sequences. Computer analysis of the TOLL sequences can be used to predict primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the TOLL sequences will yield an amplified fragment.

Somatic cell hybrids are prepared by fusing somatic cells from different mammals (e.g., human and mouse cells). As hybrids of human and mouse cells grow and divide, they gradually lose human chromosomes in random order, but retain the mouse chromosomes. By using media in which mouse cells cannot grow, because they lack a particular enzyme, but human cells can, the one human chromosome that contains the gene encoding the needed enzyme, will be retained. By using various media, panels of hybrid cell lines can be established. Each cell line in a panel contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, allowing easy mapping of individual genes to specific human chromosomes. (D'Eustachio P. et al. (1983) *Science* 220:919-924). Somatic cell hybrids containing only fragments of human chromosomes can also be produced by using human chromosomes with translocations and deletions.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be assigned per day using a single thermal cycler. Using the TOLL nucleotide sequences to design oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes. Other mapping strategies which can similarly be used to map a TOLL sequence to its chromosome include *in situ* hybridization (described in Fan, Y. et al. (1990) *Proc. Natl. Acad. Sci. USA*, 87:6223-27), pre-screening with labeled flow-sorted chromosomes, and pre-selection by hybridization to chromosome specific cDNA libraries.

5 Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase  
chromosomal spread can further be used to provide a precise chromosomal location in  
one step. Chromosome spreads can be made using cells whose division has been  
10 blocked in metaphase by a chemical such as colcemid that disrupts the mitotic spindle.

5 The chromosomes can be treated briefly with trypsin, and then stained with Giemsa. A  
pattern of light and dark bands develops on each chromosome, so that the chromosomes  
can be identified individually. The FISH technique can be used with a DNA sequence  
15 as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher  
likelihood of binding to a unique chromosomal location with sufficient signal intensity  
for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases will  
20 suffice to get good results at a reasonable amount of time. For a review of this  
technique, see Verma et al., Human Chromosomes: A Manual of Basic Techniques  
(Pergamon Press, New York 1988).

25 Reagents for chromosome mapping can be used individually to mark a single  
15 chromosome or a single site on that chromosome, or panels of reagents can be used for  
marking multiple sites and/or multiple chromosomes. Reagents corresponding to  
noncoding regions of the genes actually are preferred for mapping purposes. Coding  
30 sequences are more likely to be conserved within gene families, thus increasing the  
chance of cross hybridizations during chromosomal mapping.

20 Once a sequence has been mapped to a precise chromosomal location, the  
physical position of the sequence on the chromosome can be correlated with genetic map  
35 data. (Such data are found, for example, in V. McKusick, Mendelian Inheritance in  
Man, available on-line through Johns Hopkins University Welch Medical Library). The  
relationship between a gene and a disease, mapped to the same chromosomal region, can  
40 then be identified through linkage analysis (co-inheritance of physically adjacent genes),  
25 described in, for example, Egeland, J. et al. (1987) *Nature*, 325:783-787.

Moreover, differences in the DNA sequences between individuals affected and  
45 unaffected with a disease associated with the TOLL gene, can be determined. If a  
mutation is observed in some or all of the affected individuals but not in any unaffected  
30 individuals, then the mutation is likely to be the causative agent of the particular disease.  
50 Comparison of affected and unaffected individuals generally involves first looking for

5 structural alterations in the chromosomes, such as deletions or translocations that are  
visible from chromosome spreads or detectable using PCR based on that DNA sequence.  
10 Ultimately, complete sequencing of genes from several individuals can be performed to  
confirm the presence of a mutation and to distinguish mutations from polymorphisms.

## 2. Tissue Typing

15 The TOLL sequences of the present invention can also be used to identify  
individuals from minute biological samples. The United States military, for example, is  
considering the use of restriction fragment length polymorphism (RFLP) for  
20 identification of its personnel. In this technique, an individual's genomic DNA is  
digested with one or more restriction enzymes, and probed on a Southern blot to yield  
unique bands for identification. This method does not suffer from the current limitations  
of "Dog Tags" which can be lost, switched, or stolen, making positive identification  
25 difficult. The sequences of the present invention are useful as additional DNA markers  
for RFLP (described in U.S. Patent 5,272,057).

Furthermore, the sequences of the present invention can be used to provide an  
30 alternative technique which determines the actual base-by-base DNA sequence of  
selected portions of an individual's genome. Thus, the TOLL nucleotide sequences  
described herein can be used to prepare two PCR primers from the 5' and 3' ends of the  
20 sequences. These primers can then be used to amplify an individual's DNA and  
subsequently sequence it.

35 Panels of corresponding DNA sequences from individuals, prepared in this  
manner, can provide unique individual identifications, as each individual will have a  
unique set of such DNA sequences due to allelic differences. The sequences of the  
40 present invention can be used to obtain such identification sequences from individuals  
and from tissue. The TOLL nucleotide sequences of the invention uniquely represent  
portions of the human genome. Allelic variation occurs to some degree in the coding  
45 regions of these sequences, and to a greater degree in the noncoding regions. It is  
estimated that allelic variation between individual humans occurs with a frequency of  
30 about once per each 500 bases. Each of the sequences described herein can, to some  
degree, be used as a standard against which DNA from an individual can be compared  
50

5 for identification purposes. Because greater numbers of polymorphisms occur in the  
noncoding regions, fewer sequences are necessary to differentiate individuals. The  
noncoding sequences of SEQ ID NO:1 can comfortably provide positive individual  
10 identification with a panel of perhaps 10 to 1,000 primers which each yield a noncoding  
5 amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ  
ID NO:3 are used, a more appropriate number of primers for positive individual  
15 identification would be 500-2,000.

If a panel of reagents from TOLL nucleotide sequences described herein is used  
to generate a unique identification database for an individual, those same reagents can  
10 later be used to identify tissue from that individual. Using the unique identification  
20 database, positive identification of the individual, living or dead, can be made from  
extremely small tissue samples.

### 25 3. Use of Partial TOLL Sequences in Forensic Biology

15 DNA-based identification techniques can also be used in forensic biology.  
Forensic biology is a scientific field employing genetic typing of biological evidence  
found at a crime scene as a means for positively identifying, for example, a perpetrator  
30 of a crime. To make such an identification, PCR technology can be used to amplify  
DNA sequences taken from very small biological samples such as tissues, e.g., hair or  
20 skin, or body fluids, e.g., blood, saliva, or semen found at a crime scene. The amplified  
35 sequence can then be compared to a standard, thereby allowing identification of the  
origin of the biological sample.

The sequences of the present invention can be used to provide polynucleotide  
reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can  
40 enhance the reliability of DNA-based forensic identifications by, for example, providing  
25 another "identification marker" (i.e. another DNA sequence that is unique to a particular  
individual). As mentioned above, actual base sequence information can be used for  
45 identification as an accurate alternative to patterns formed by restriction enzyme  
generated fragments. Sequences targeted to noncoding regions of SEQ ID NO:1 are  
30 particularly appropriate for this use as greater numbers of polymorphisms occur in the  
noncoding regions, making it easier to differentiate individuals using this technique.  
50

5 Examples of polynucleotide reagents include the TOLL nucleotide sequences or portions thereof, e.g., fragments derived from the noncoding regions of SEQ ID NO:1 having a length of at least 20 bases, preferably at least 30 bases.

10 The TOLL nucleotide sequences described herein can further be used to provide  
5 polynucleotide reagents, e.g., labeled or labelable probes which can be used in, for example, an *in situ* hybridization technique, to identify a specific tissue, e.g., brain tissue. This can be very useful in cases where a forensic pathologist is presented with a tissue of unknown origin. Panels of such TOLL probes can be used to identify tissue by species and/or by organ type.

15 In a similar fashion, these reagents, e.g., TOLL primers or probes can be used to  
20 screen tissue culture for contamination (i.e. screen for the presence of a mixture of different types of cells in a culture).

### 25 C. Predictive Medicine:

15 The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically.  
30 Accordingly, one aspect of the present invention relates to diagnostic assays for determining TOLL protein and/or nucleic acid expression as well as TOLL activity, in the context of a biological sample (e.g., blood, serum, cells, tissue) to thereby determine  
20 whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant or unwanted TOLL expression or activity. The invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with TOLL protein, nucleic acid  
35 expression or activity. For example, mutations in a TOLL gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with TOLL protein, nucleic acid expression or activity.

40 Another aspect of the invention pertains to monitoring the influence of agents  
30 (e.g., drugs, compounds) on the expression or activity of TOLL in clinical trials.

50 These and other agents are described in further detail in the following sections.



5                   1.     Diagnostic Assays

                  An exemplary method for detecting the presence or absence of TOLL protein or  
                  nucleic acid in a biological sample involves obtaining a biological sample from a test  
10               subject and contacting the biological sample with a compound or an agent capable of  
                  5     detecting TOLL protein or nucleic acid (e.g., mRNA, genomic DNA) that encodes  
                  TOLL protein such that the presence of TOLL protein or nucleic acid is detected in the  
                  biological sample. A preferred agent for detecting TOLL mRNA or genomic DNA is a  
15               labeled nucleic acid probe capable of hybridizing to TOLL mRNA or genomic DNA.  
                  The nucleic acid probe can be, for example, a full-length TOLL nucleic acid, such as the  
                  10     nucleic acid of SEQ ID NO:1, 3, or 4, or the DNA insert of the plasmid deposited with  
                  ATCC as Accession Number \_\_\_\_\_, or a portion thereof, such as an oligonucleotide of  
20               at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically  
                  hybridize under stringent conditions to TOLL mRNA or genomic DNA. Other suitable  
                  25     probes for use in the diagnostic assays of the invention are described herein.

15               A preferred agent for detecting TOLL protein is an antibody capable of binding  
                  to TOLL protein, preferably an antibody with a detectable label. Antibodies can be  
                  polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof  
30               (e.g., Fab or F(ab')<sub>2</sub>) can be used. The term "labeled", with regard to the probe or  
                  antibody, is intended to encompass direct labeling of the probe or antibody by coupling  
20               (i.e., physically linking) a detectable substance to the probe or antibody, as well as  
                  indirect labeling of the probe or antibody by reactivity with another reagent that is  
35               directly labeled. Examples of indirect labeling include detection of a primary antibody  
                  using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with  
                  biotin such that it can be detected with fluorescently labeled streptavidin. The term  
40               25     "biological sample" is intended to include tissues, cells and biological fluids isolated  
                  from a subject, as well as tissues, cells and fluids present within a subject. That is, the  
                  detection method of the invention can be used to detect TOLL mRNA, protein, or  
45               genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro*  
                  techniques for detection of TOLL mRNA include Northern hybridizations and *in situ*  
                  30     hybridizations. *In vitro* techniques for detection of TOLL protein include enzyme linked  
                  immunosorbent assays (ELISAs), Western blots, immunoprecipitations and  
50

5 immunofluorescence. *In vitro* techniques for detection of TOLL genomic DNA include  
Southern hybridizations. Furthermore, *in vivo* techniques for detection of TOLL protein  
10 include introducing into a subject a labeled anti-TOLL antibody. For example, the  
antibody can be labeled with a radioactive marker whose presence and location in a  
5 subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the  
15 test subject. Alternatively, the biological sample can contain mRNA molecules from the  
test subject or genomic DNA molecules from the test subject. A preferred biological  
sample is a serum sample isolated by conventional means from a subject.

10 In another embodiment, the methods further involve obtaining a control  
biological sample from a control subject, contacting the control sample with a  
20 compound or agent capable of detecting TOLL protein, mRNA, or genomic DNA, such  
that the presence of TOLL protein, mRNA or genomic DNA is detected in the biological  
sample, and comparing the presence of TOLL protein, mRNA or genomic DNA in the  
25 control sample with the presence of TOLL protein, mRNA or genomic DNA in the test  
15 sample.

The invention also encompasses kits for detecting the presence of TOLL in a  
30 biological sample. For example, the kit can comprise a labeled compound or agent  
capable of detecting TOLL protein or mRNA in a biological sample; means for  
20 determining the amount of TOLL in the sample; and means for comparing the amount of  
TOLL in the sample with a standard. The compound or agent can be packaged in a  
35 suitable container. The kit can further comprise instructions for using the kit to detect  
TOLL protein or nucleic acid.

## 40 2. Prognostic Assays

25 The diagnostic methods described herein can furthermore be utilized to identify  
subjects having or at risk of developing a disease or disorder associated with aberrant or  
unwanted TOLL expression or activity. As used herein, the term "aberrant" includes a  
45 TOLL expression or activity which deviates from the wild type TOLL expression or  
activity. Aberrant expression or activity includes increased or decreased expression or  
30 activity, as well as expression or activity which does not follow the wild type  
developmental pattern of expression or the subcellular pattern of expression. For  
50

5 example, aberrant TOLL expression or activity is intended to include the cases in which  
a mutation in the TOLL gene causes the TOLL gene to be under-expressed or over-  
10 expressed and situations in which such mutations result in a non-functional TOLL  
protein or a protein which does not function in a wild-type fashion, e.g., a protein which  
5 does not interact with a TOLL ligand or one which interacts with a non-TOLL ligand.  
As used herein, the term "unwanted" includes an unwanted phenomenon involving a  
TOLL associated response, such as an immune response. For example, the term  
15 unwanted includes a TOLL expression or activity which is undesirable in a subject.

The assays described herein, such as the preceding diagnostic assays or the  
10 following assays, can be utilized to identify a subject having or at risk of developing a  
disorder associated with a misregulation in TOLL protein activity or nucleic acid  
20 expression, such as, for example, an immune disorder. Alternatively, the prognostic  
assays can be utilized to identify a subject having or at risk for developing a disorder  
associated with a misregulation in TOLL protein activity or nucleic acid expression.  
25 Thus, the present invention provides a method for identifying a disease or disorder  
associated with aberrant or unwanted TOLL expression or activity in which a test  
sample is obtained from a subject and TOLL protein or nucleic acid (e.g., mRNA or  
30 genomic DNA) is detected, wherein the presence of TOLL protein or nucleic acid is  
diagnostic for a subject having or at risk of developing a disease or disorder associated  
20 with aberrant or unwanted TOLL expression or activity. As used herein, a "test sample"  
refers to a biological sample obtained from a subject of interest. For example, a test  
35 sample can be a biological fluid (e.g., serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine  
whether a subject can be administered an agent (e.g., an agonist, antagonist,  
40 25 peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate)  
to treat a disease or disorder associated with aberrant or unwanted TOLL expression or  
activity. For example, such methods can be used to determine whether a subject can be  
45 effectively treated with an agent for a TOLL associated disorder. Thus, the present  
invention provides methods for determining whether a subject can be effectively treated  
30 with an agent for a disorder associated with aberrant or unwanted TOLL expression or  
activity in which a test sample is obtained and TOLL protein or nucleic acid expression  
50

5 or activity is detected (e.g., wherein the abundance of TOLL protein or nucleic acid expression or activity is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant or unwanted TOLL expression or activity).

10 The methods of the invention can also be used to detect genetic alterations in a TOLL gene, thereby determining if a subject with the altered gene is at risk for a disorder characterized by misregulation in TOLL protein activity or nucleic acid expression. In preferred embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic alteration characterized by at least one of an alteration affecting the integrity of a gene encoding a TOLL-protein, or 15 the mis-expression of the TOLL gene. For example, such genetic alterations can be detected by ascertaining the existence of at least one of 1) a deletion of one or more nucleotides from a TOLL gene; 2) an addition of one or more nucleotides to a TOLL gene; 3) a substitution of one or more nucleotides of a TOLL gene, 4) a chromosomal rearrangement of a TOLL gene; 5) an alteration in the level of a messenger RNA 20 transcript of a TOLL gene, 6) aberrant modification of a TOLL gene, such as of the methylation pattern of the genomic DNA, 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of a TOLL gene, 8) a non-wild type level of a TOLL-protein, 9) allelic loss of a TOLL gene, and 10) inappropriate post-translational modification of a TOLL-protein. As described herein, there are a large number of assays 25 known in the art which can be used for detecting alterations in a TOLL gene. A preferred biological sample is a tissue or serum sample isolated by conventional means from a subject.

In certain embodiments, detection of the alteration involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g., U.S. Patent Nos. 40 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran et al. (1988) *Science* 241:1077-1080; and Nakazawa et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:360-364), the latter of which can be particularly useful for detecting point mutations in the TOLL gene (see Abravaya et al. (1995) *Nucleic Acids Res.* 23:675-682). This method can include the steps of 45 collecting a sample of cells from a subject, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or 50

5 more primers which specifically hybridize to a TOLL gene under conditions such that  
hybridization and amplification of the TOLL gene (if present) occurs, and detecting the  
10 presence or absence of an amplification product, or detecting the size of the  
amplification product and comparing the length to a control sample. It is anticipated  
5 that PCR and/or LCR may be desirable to use as a preliminary amplification step in  
conjunction with any of the techniques used for detecting mutations described herein.

15 Alternative amplification methods include: self sustained sequence replication  
(Guatelli, J.C. et al., (1990) *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional  
amplification system (Kwoh, D.Y. et al., (1989) *Proc. Natl. Acad. Sci. USA* 86:1173-  
10 1177), Q-Beta Replicase (Lizardi, P.M. et al. (1988) *Bio-Technology* 6:1197), or any  
20 other nucleic acid amplification method, followed by the detection of the amplified  
molecules using techniques well known to those of skill in the art. These detection  
schemes are especially useful for the detection of nucleic acid molecules if such  
25 molecules are present in very low numbers.

15 In an alternative embodiment, mutations in a TOLL gene from a sample cell can  
be identified by alterations in restriction enzyme cleavage patterns. For example,  
sample and control DNA is isolated, amplified (optionally), digested with one or more  
30 restriction endonucleases, and fragment length sizes are determined by gel  
electrophoresis and compared. Differences in fragment length sizes between sample and  
20 control DNA indicates mutations in the sample DNA. Moreover, the use of sequence  
specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score  
35 for the presence of specific mutations by development or loss of a ribozyme cleavage  
site.

40 In other embodiments, genetic mutations in TOLL can be identified by  
25 hybridizing a sample and control nucleic acids, e.g., DNA or RNA, to high density  
arrays containing hundreds or thousands of oligonucleotides probes (Cronin, M.T. et al.  
(1996) *Human Mutation* 7: 244-255; Kozal, M.J. et al. (1996) *Nature Medicine* 2: 753-  
45 759). For example, genetic mutations in TOLL can be identified in two dimensional  
arrays containing light-generated DNA probes as described in Cronin, M.T. *et al. supra*.  
30 Briefly, a first hybridization array of probes can be used to scan through long stretches  
50 of DNA in a sample and control to identify base changes between the sequences by

5 making linear arrays of sequential overlapping probes. This step allows the  
identification of point mutations. This step is followed by a second hybridization array  
that allows the characterization of specific mutations by using smaller, specialized probe  
10 arrays complementary to all variants or mutations detected. Each mutation array is  
5 composed of parallel probe sets, one complementary to the wild-type gene and the other  
complementary to the mutant gene.

15 In yet another embodiment, any of a variety of sequencing reactions known in  
the art can be used to directly sequence the TOLL gene and detect mutations by  
comparing the sequence of the sample TOLL with the corresponding wild-type (control)  
10 sequence. Examples of sequencing reactions include those based on techniques  
developed by Maxam and Gilbert ((1977) *Proc. Natl. Acad. Sci. USA* 74:560) or Sanger  
20 ((1977) *Proc. Natl. Acad. Sci. USA* 74:5463). It is also contemplated that any of a  
variety of automated sequencing procedures can be utilized when performing the  
diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass  
25 spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen et al.  
15 (1996) *Adv. Chromatogr.* 36:127-162; and Griffin et al. (1993) *Appl. Biochem.*  
*Biotechnol.* 38:147-159).

30 Other methods for detecting mutations in the TOLL gene include methods in  
which protection from cleavage agents is used to detect mismatched bases in RNA/RNA  
20 or RNA/DNA heteroduplexes (Myers et al. (1985) *Science* 230:1242). In general, the art  
technique of "mismatch cleavage" starts by providing heteroduplexes of formed by  
35 hybridizing (labeled) RNA or DNA containing the wild-type TOLL sequence with  
potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded  
duplexes are treated with an agent which cleaves single-stranded regions of the duplex  
40 such as which will exist due to basepair mismatches between the control and sample  
25 strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA  
hybrids treated with S1 nuclease to enzymatically digesting the mismatched regions. In  
45 other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with  
hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched  
30 regions. After digestion of the mismatched regions, the resulting material is then  
50 separated by size on denaturing polyacrylamide gels to determine the site of mutation.

5 See, for example, Cotton *et al.* (1988) *Proc. Natl Acad Sci USA* 85:4397; Saleeba *et al.* (1992) *Methods Enzymol.* 217:286-295. In a preferred embodiment, the control DNA or RNA can be labeled for detection.

10 In still another embodiment, the mismatch cleavage reaction employs one or  
5 more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in TOLL cDNAs obtained from samples of cells. For example, the mutY  
15 enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu *et al.* (1994) *Carcinogenesis*  
20 15:1657-1662). According to an exemplary embodiment, a probe based on a TOLL sequence, e.g., a wild-type TOLL sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the  
25 like. See, for example, U.S. Patent No. 5,459,039.

15 In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in TOLL genes. For example, single strand conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility  
30 between mutant and wild type nucleic acids (orita *et al.* (1989) *Proc Natl. Acad. Sci USA*: 86:2766. see also Cotton (1993) *Mutat. Res.* 285:125-144; and Hayashi (1992) *Genet. Anal. Tech. Appl.* 9:73-79). Single-stranded DNA fragments of sample and control TOLL nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the resulting  
35 alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of  
40 the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a preferred embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex  
45 molecules on the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet* 7:5).

5 In yet another embodiment the movement of mutant or wild-type fragments in  
polyacrylamide gels containing a gradient of denaturant is assayed using denaturing  
10 gradient gel electrophoresis (DGGE) (Myers *et al.* (1985) *Nature* 313:495). When  
DGGE is used as the method of analysis, DNA will be modified to insure that it does not  
5 completely denature, for example by adding a GC clamp of approximately 40 bp of  
high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is  
15 used in place of a denaturing gradient to identify differences in the mobility of control  
and sample DNA (Rosenbaum and Reissner (1987) *Biophys Chem* 265:12753).

Examples of other techniques for detecting point mutations include, but are not  
10 limited to, selective oligonucleotide hybridization, selective amplification, or selective  
primer extension. For example, oligonucleotide primers may be prepared in which the  
20 known mutation is placed centrally and then hybridized to target DNA under conditions  
which permit hybridization only if a perfect match is found (Saiki *et al.* (1986) *Nature*  
324:163); Saiki *et al.* (1989) *Proc. Natl Acad. Sci USA* 86:6230). Such allele specific  
25 oligonucleotides are hybridized to PCR amplified target DNA or a number of different  
mutations when the oligonucleotides are attached to the hybridizing membrane and  
30 hybridized with labeled target DNA.

Alternatively, allele specific amplification technology which depends on selective  
PCR amplification may be used in conjunction with the instant invention.

20 Oligonucleotides used as primers for specific amplification may carry the mutation of  
interest in the center of the molecule (so that amplification depends on differential  
35 hybridization) (Gibbs *et al.* (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme 3'  
end of one primer where, under appropriate conditions, mismatch can prevent, or reduce  
40 polymerase extension (Prossner (1993) *Tibtech* 11:238). In addition it may be desirable  
25 to introduce a novel restriction site in the region of the mutation to create cleavage-based  
detection (Gasparini *et al.* (1992) *Mol. Cell Probes* 6:1). It is anticipated that in certain  
embodiments amplification may also be performed using Taq ligase for amplification  
45 (Barany (1991) *Proc. Natl. Acad. Sci USA* 88:189). In such cases, ligation will occur  
only if there is a perfect match at the 3' end of the 5' sequence making it possible to detect  
30 the presence of a known mutation at a specific site by looking for the presence or absence  
of amplification.  
50



5 The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.*, in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving a TOLL gene.

Furthermore, any cell type or tissue in which TOLL is expressed may be utilized in the prognostic assays described herein.

### 3. Monitoring of Effects During Clinical Trials

10 Monitoring the influence of agents (*e.g.*, drugs) on the expression or activity of a TOLL protein (*e.g.*, the modulation of signaling mechanisms involved in immunity) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase TOLL gene expression, protein levels, or upregulate TOLL activity, can be monitored in clinical trials of subjects exhibiting decreased TOLL gene expression, protein levels, or downregulated TOLL activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease TOLL gene expression, protein levels, or downregulate TOLL activity, can be monitored in clinical trials of subjects exhibiting increased TOLL gene expression, protein levels, or upregulated TOLL activity. In such clinical trials, the expression or activity of a TOLL gene, and preferably, other genes that have been implicated in, for example, a TOLL-associated disorder can be used as a "read out" or markers of the phenotype of a particular cell.

For example, and not by way of limitation, genes, including TOLL, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule) which modulates TOLL activity (*e.g.*, identified in a screening assay as described herein) can be identified. Thus, to study the effect of agents on TOLL-associated disorders (*e.g.*, disorders characterized by deregulated signaling mechanisms involved in immunity), for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of TOLL and other genes implicated in the TOLL-associated disorder, respectively. The levels of gene expression (*e.g.*, a gene expression pattern) can be quantified by northern blot analysis or RT-PCR, as described herein, or

alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of TOLL or other genes. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during treatment of the individual with the agent.

In a preferred embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (e.g., an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) including the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of a TOLL protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the TOLL protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the TOLL protein, mRNA, or genomic DNA in the pre-administration sample with the TOLL protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of TOLL to higher levels than detected, i.e., to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of TOLL to lower levels than detected, i.e. to decrease the effectiveness of the agent. According to such an embodiment, TOLL expression or activity may be used as an indicator of the effectiveness of an agent, even in the absence of an observable phenotypic response.

#### D. Methods of Treatment:

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant or unwanted TOLL expression or activity. With regards to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics.

5 "Pharmacogenomics", as used herein, refers to the application of genomics technologies  
such as gene sequencing, statistical genetics, and gene expression analysis to drugs in  
clinical development and on the market. More specifically, the term refers the study of  
10 how a patient's genes determine his or her response to a drug (e.g., a patient's "drug  
5 response phenotype", or "drug response genotype".) Thus, another aspect of the  
invention provides methods for tailoring an individual's prophylactic or therapeutic  
treatment with either the TOLL molecules of the present invention or TOLL modulators  
15 according to that individual's drug response genotype. Pharmacogenomics allows a  
clinician or physician to target prophylactic or therapeutic treatments to patients who  
20 will most benefit from the treatment and to avoid treatment of patients who will  
experience toxic drug-related side effects.

#### 1. Prophylactic Methods

25 In one aspect, the invention provides a method for preventing in a subject, a  
15 disease or condition associated with an aberrant or unwanted TOLL expression or  
activity, by administering to the subject a TOLL protein or an agent which modulates  
TOLL expression or at least one TOLL activity. Subjects at risk for a disease which is  
30 caused or contributed to by aberrant or unwanted TOLL expression or activity can be  
identified by, for example, any or a combination of diagnostic or prognostic assays as  
20 described herein. Administration of a prophylactic agent can occur prior to the  
manifestation of symptoms characteristic of the TOLL aberrancy, such that a disease or  
35 disorder is prevented or, alternatively, delayed in its progression. Depending on the type  
of TOLL aberrancy, for example, a TOLL protein, TOLL agonist or TOLL antagonist  
agent can be used for treating the subject. The appropriate agent can be determined  
40 based on screening assays described herein.  
25

#### 2. Therapeutic Methods

45 Another aspect of the invention pertains to methods of modulating TOLL  
expression or activity for therapeutic purposes. Accordingly, in an exemplary  
30 embodiment, the modulatory method of the invention involves contacting a cell with a  
TOLL protein or agent that modulates one or more of the activities of TOLL protein  
50

5 activity associated with the cell. An agent that modulates TOLL protein activity can be  
an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring  
10 target molecule of a TOLL protein (e.g., a TOLL substrate), a TOLL antibody, a TOLL  
agonist or antagonist, a peptidomimetic of a TOLL agonist or antagonist, or other small  
5 molecule. In one embodiment, the agent stimulates one or more TOLL activities.  
Examples of such stimulatory agents include active TOLL protein and a nucleic acid  
15 molecule encoding TOLL that has been introduced into the cell. In another  
embodiment, the agent inhibits one or more TOLL activities. Examples of such  
inhibitory agents include antisense TOLL nucleic acid molecules, anti-TOLL antibodies,  
20 and TOLL inhibitors. These modulatory methods can be performed *in vitro* (e.g., by  
culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the  
agent to a subject). As such, the present invention provides methods of treating an  
individual afflicted with a disease or disorder characterized by aberrant or unwanted  
25 expression or activity of a TOLL protein or nucleic acid molecule (e.g., rheumatoid  
arthritis, systemic lupus erythematosus, myasthenia gravis, Grave's disease, Sjogren  
15 syndrome, polymyositis and dermatomyositis, psoriasis, pemphigus vulgaris, bullous  
pemphigoid, inflammatory bowel disease, Kawasaki disease, asthma, and graft vs. host  
30 disease). In one embodiment, the method involves administering an agent (e.g., an agent  
identified by a screening assay described herein), or combination of agents that  
20 modulates (e.g., upregulates or downregulates) TOLL expression or activity. In another  
embodiment, the method involves administering a TOLL protein or nucleic acid  
35 molecule as therapy to compensate for reduced, aberrant, or unwanted TOLL expression  
or activity.

40 Stimulation of TOLL activity is desirable in situations in which TOLL is  
25 abnormally downregulated and/or in which increased TOLL activity is likely to have a  
beneficial effect. For example, stimulation of TOLL activity is desirable in situations in  
which a TOLL is downregulated and/or in which increased TOLL activity is likely to  
45 have a beneficial effect. Likewise, inhibition of TOLL activity is desirable in situations  
in which TOLL is abnormally upregulated and/or in which decreased TOLL activity is  
30 likely to have a beneficial effect.

### 3. Pharmacogenomics

The TOLL molecules of the present invention, as well as agents, or modulators which have a stimulatory or inhibitory effect on TOLL activity (e.g., TOLL gene expression) as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) TOLL-associated disorders associated with aberrant or unwanted TOLL activity. In conjunction with such treatment, pharmacogenomics (i.e., the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer a TOLL molecule or TOLL modulator as well as tailoring the dosage and/or therapeutic regimen of treatment with a TOLL molecule or TOLL modulator.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, for example, Eichelbaum, M. et al. (1996) *Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 and Linder, M.W. et al. (1997) *Clin. Chem.* 43(2):254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare genetic defects or as naturally-occurring polymorphisms. For example, glucose-6-phosphate dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

One pharmacogenomics approach to identifying genes that predict drug response, known as "a genome-wide association", relies primarily on a high-resolution map of the human genome consisting of already known gene-related markers (e.g., a "bi-allelic" gene marker map which consists of 60,000-100,000 polymorphic or variable

5 sites on the human genome, each of which has two variants.) Such a high-resolution genetic map can be compared to a map of the genome of each of a statistically significant number of patients taking part in a Phase II/III drug trial to identify markers associated with a particular observed drug response or side effect. Alternatively, such a high resolution map can be generated from a combination of some ten-million known single nucleotide polymorphisms (SNPs) in the human genome. As used herein, a "SNP" is a common alteration that occurs in a single nucleotide base in a stretch of DNA. For example, a SNP may occur once per every 1000 bases of DNA. A SNP may be involved in a disease process, however, the vast majority may not be disease-associated. Given a genetic map based on the occurrence of such SNPs, individuals can be grouped into genetic categories depending on a particular pattern of SNPs in their individual genome. In such a manner, treatment regimens can be tailored to groups of genetically similar individuals, taking into account traits that may be common among such genetically similar individuals.

15 Alternatively, a method termed the "candidate gene approach", can be utilized to identify genes that predict drug response. According to this method, if a gene that encodes a drug target is known (e.g., a TOLL protein of the present invention), all common variants of that gene can be fairly easily identified in the population and it can be determined if having one version of the gene versus another is associated with a particular drug response.

35 As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (e.g., N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently

5 experience exaggerated drug response and side effects when they receive standard doses.

10 If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. The other extreme are the so called ultra-rapid metabolizers who  
5 do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

15 Alternatively, a method termed the "gene expression profiling", can be utilized to identify genes that predict drug response. For example, the gene expression of an animal dosed with a drug (e.g., a TOLL molecule or TOLL modulator of the present  
10 invention) can give an indication whether gene pathways related to toxicity have been turned on.

20 Information generated from more than one of the above pharmacogenomics approaches can be used to determine appropriate dosage and treatment regimens for prophylactic or therapeutic treatment an individual. This knowledge, when applied to  
25 dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a TOLL molecule or TOLL modulator, such as a modulator identified by one of the exemplary  
30 screening assays described herein.

35 This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are incorporated herein by reference.

## EXAMPLES

### 40 25 EXAMPLE 1: IDENTIFICATION AND CHARACTERIZATION OF HUMAN AND MOUSE TOLL cDNA

45 In this example, the identification and characterization of the gene encoding human TOLL (clone jthKa089g09) and mouse TOLL (clone jtmBa212b08) is described.

30

50

55

Isolation of the Human TOLL cDNA

The invention is based, at least in part, on the discovery of a human gene encoding a novel protein, referred to herein as TOLL. A clone was originally identified based on sequence homology to Toll proteins, and based on the sequence of this first clone, primers were designed and used to screen a human fat cell library (obtained from Clontech). A positive human clone, jthKa089g09, was identified. The entire sequence of the human clone was determined and found to contain an open reading frame termed human "TOLL".

The nucleotide sequence encoding the human TOLL protein is shown in Figure 1 and is set forth as SEQ ID NO:1. The full length protein encoded by this nucleic acid comprises about 548 amino acids and has the amino acid sequence shown in Figure 1 and set forth as SEQ ID NO:2. The coding region (open reading frame) of SEQ ID NO:1 is set forth as SEQ ID NO:3. Clone jthKa089g09, comprising the entire coding region of the human TOLL gene, was deposited with the American Type Culture Collection (ATCC®), 10801 University Boulevard, Manassas, VA 20110-2209, on \_\_\_\_\_, and assigned Accession No. \_\_\_\_\_.

Isolation of the Mouse TOLL cDNA

The invention is further based, at least in part, on the discovery of a partial mouse nucleic acid sequence, referred to herein as mouse TOLL nucleic acid molecule. A clone was originally identified based on sequence homology to Toll proteins, and based on the sequence of this first clone, primers were designed and used to screen a mouse cell library. A positive mouse clone, jtmBa212b08, was identified. The sequence of this mouse clone was determined and is termed mouse "TOLL".

A partial nucleotide sequence of the mouse TOLL gene is shown in Figure 3 and is set forth as SEQ ID NO:4. Clone jtmBa212b08, comprising the entire sequence of SEQ ID NO:4, was deposited with the American Type Culture Collection (ATCC®), 10801 University Boulevard, Manassas, VA 20110-2209, on \_\_\_\_\_, and assigned Accession No. \_\_\_\_\_.



Analysis of the Human TOLL Molecule

A BLASTX 1.4.9 search, using a score of 100 and a word length of 3 (Altschul et al. (1990) *J. Mol. Biol.* 215:403) of the translated nucleotide sequence of human TOLL revealed that human TOLL is similar to the rat MEGF5 protein (Accession Number AB011531), the neurogenic extracellular slit protein slit2 (Accession Number AF055585), human slit-2 protein (Accession Number AB017168), human slit-3 protein (Accession Number AB017169), human slit-1 protein (Accession Number AB017167) and the rat MEGF4 protein (Accession Number AB014462). The human TOLL protein is 39% identical to the rat MEGF5 protein (Accession Number AB011531) over translated nucleotides 191 to 643, 30% identical to this sequence over translated nucleotides 188 to 715, 28% identical to this sequence over translated nucleotides 347 to 787, 30% identical to this sequence over translated nucleotides 842 to 1189, 39% identical to this sequence over translated nucleotides 914 to 1189, 27% identical to this sequence over translated nucleotides 842 to 1276, 35% identical over translated nucleotides 320 to 643, 37% identical over translated nucleotides 482 to 715, and 36% identical over translated nucleotides 839 to 946. The human TOLL protein is 37% identical to the neurogenic extracellular slit protein slit2 (Accession Number AF055585) over translated nucleotides 191 to 643, 30% identical to this sequence over translated nucleotides 188 to 730, 38% identical to this sequence over translated nucleotides 911 to 1189, 28% identical to this sequence over translated nucleotides 842 to 1246, 26% identical to this sequence over translated nucleotides 347 to 775, 36% identical to this sequence over translated nucleotides 311 to 586, 30% identical to this sequence over translated nucleotides 338 to 715, 28% identical to this sequence over residues 842 to 1258, 26% identical to this sequence over translated nucleotides 410 to 805, 22% identical to this sequence over translated nucleotides 842 to 1276, 31% identical to this sequence over translated nucleotides 320 to 643, 25% identical to this sequence over translated nucleotides 275 to 601, 35% identical to this sequence over translated nucleotides 272 to 514, 29% identical to this sequence over translated nucleotides 419 to 790, 27% identical to this sequence over translated nucleotides 386 to 715, 30% identical to this sequence over translated nucleotides 914 to 1168, 28% identical to this sequence over translated nucleotides 839 to 1156, 28% identical to this sequence over

5 translated nucleotides 275 to 571, 28% identical to this sequence over translated  
nucleotides 851 to 1168, 27% identical to this sequence over translated nucleotides 635  
10 to 1090, 29% identical to this sequence over translated nucleotides 491 to 793, 26%  
identical to this sequence over translated nucleotides 263 to 601, is 27% identical to this  
5 sequence over translated nucleotides 458 to 787, 26% identical to this sequence over  
translated nucleotides 842 to 1162, 23% identical to this sequence over translated  
15 nucleotides 695 to 1084, 35% identical to this sequence over translated nucleotides 479  
to 709, 30% identical to this sequence over translated nucleotides 551 to 802, 25%  
identical to this sequence over translated nucleotides 896 to 1162, 38% identical to this  
20 sequence over translated nucleotides 275 to 436, 25% identical to this sequence over  
translated nucleotides 383 to 643, 28% identical to this sequence over translated  
nucleotides 839 to 1090, and 30% identical to this sequence over translated nucleotides  
281 to 508.

25 A BLASTN 1.4.9 search, using a score of 100 and a word length of 12 (Altschul  
15 et al. (1990) *J. Mol. Biol.* 215:403) of the nucleotide sequence of human TOLL revealed  
that the TOLL sequence is similar to the mouse mRNA for Knowles Solter mouse 2 cell  
30 *Mus musculus* cDNA clone 1124974 (Accession Number AA692768), to Soares infant  
brain 1NIB Homo sapiens cDNA clone mRNA (Accession Number R54798), to Soares  
infant brain 1NIB Homo sapiens cDNA clone mRNA (Accession Number H12046) and  
20 to Soares infant brain 1NIB Homo sapiens cDNA clone mRNA (Accession Number  
35 H12045). The TOLL nucleic acid molecule is 92% identical to the mouse mRNA for  
Knowles Solter mouse 2 cell *Mus musculus* cDNA clone 1124974 (Accession Number  
AA692768) over nucleotides 1038 to 548. The TOLL nucleic acid molecule is 100%  
40 identical to Soares infant brain 1NIB Homo sapiens cDNA clone mRNA (Accession  
25 Number R54798) over nucleotides 2061 to 1709, and 1711 to 1664, is 84% identical to  
this same sequence over nucleotides 1641-1603, and is 88% identical to this sequence  
over nucleotides 1656-1622. The TOLL nucleic acid molecule is 99% identical to  
45 Soares infant brain 1NIB Homo sapiens cDNA clone mRNA (Accession Number  
H12046) over nucleotides 2060 to 1850, is 98% identical to this sequence over  
30 nucleotides 1859-1709, is 75% identical to this sequence over nucleotides 1711-1658, is  
50 92% identical to this sequence over nucleotides 1629-1603 and is 62% identical to this

5 sequence over nucleotides 1685-1622. The TOLL nucleic acid molecule is 98%  
identical to Soares infant brain INIB Homo sapiens cDNA clone mRNA (Accession  
Number H12045) over nucleotides 1323-1535, is 73% identical to this sequence over  
10 nucleotides 1584 to 1680, is 67% identical to this sequence over nucleotides 1516-1614,  
5 is 100% identical to this sequence over nucleotides 1576-1615, is 94% identical to this  
sequence over nucleotides 1686 to 1719, is 82% identical to this sequence over  
15 nucleotides 1663 to 1697, and is 66% identical to this sequence over nucleotides 1582 to  
1629.

#### 10 Tissue Distribution of TOLL mRNA

20 This Example describes the tissue distribution of TOLL mRNA, as can be  
determined by Northern blot hybridization and *in situ* hybridization.

Northern blot hybridizations with the various RNA samples are performed under  
25 standard conditions and washed under stringent conditions, i.e., 0.2xSSC at 65°C. The  
15 DNA probe is radioactively labeled with <sup>32</sup>P-dCTP using the Prime-It kit (Stratagene,  
La Jolla, CA) according to the instructions of the supplier. Filters containing human or  
mouse mRNA (MultiTissue Northern I and MultiTissue Northern II from Clontech, Palo  
30 Alto, CA) are probed in ExpressHyb hybridization solution (Clontech) and washed at  
high stringency according to manufacturer's recommendations.

20 For *in situ* analysis, various tissues obtained from brains, e.g. rat or monkey  
35 brains, are first frozen on dry ice. Ten-micrometer-thick coronal sections of the tissues  
are postfixed with 4% formaldehyde in DEPC treated 1X phosphate- buffered saline at  
room temperature for 10 minutes before being rinsed twice in DEPC 1X phosphate-  
40 buffered saline and once in 0.1 M triethanolamine-HCl (pH 8.0). Following incubation  
25 in 0.25% acetic anhydride-0.1 M triethanolamine-HCl for 10 minutes, sections are  
rinsed in DEPC 2X SSC (1X SSC is 0.15M NaCl plus 0.015M sodium citrate). Tissue  
is then dehydrated through a series of ethanol washes, incubated in 100% chloroform for  
45 5 minutes, and then rinsed in 100% ethanol for 1 minute and 95% ethanol for 1 minute  
and allowed to air dry.

Hybridizations are performed with  $^{35}\text{S}$ -radiolabeled ( $5 \times 10^7$  cpm/ml) cRNA probes. Probes are incubated in the presence of a solution containing 600 mM NaCl, 10 mM Tris (pH 7.5), 1 mM EDTA, 0.01% sheared salmon sperm DNA, 0.01% yeast tRNA, 0.05% yeast total RNA type X1, 1 X Denhardt's solution, 50% formamide, 10% dextran sulfate, 100 mM dithiothreitol, 0.1% sodium dodecyl sulfate (SDS), and 0.1% sodium thiosulfate for 18 hours at 55°C.

After hybridization, slides are washed with 2 X SSC. Sections are then sequentially incubated at 37°C in TNE (a solution containing 10 mM Tris-HCl (pH 7.6), 500 mM NaCl, and 1 mM EDTA), for 10 minutes, in TNE with 10 µg of RNase A per ml for 30 minutes, and finally in TNE for 10 minutes. Slides are then rinsed with 2 X SSC at room temperature, washed with 2 X SSC at 50°C for 1 hour, washed with 0.2 X SSC at 55°C for 1 hour, and 0.2 X SSC at 60°C for 1 hour. Sections are then dehydrated rapidly through serial ethanol-0.3 M sodium acetate concentrations before being air dried and exposed to Kodak Biomax MR scientific imaging film for 24 hours and subsequently dipped in NB-2 photoemulsion and exposed at 4°C for 7 days before being developed and counter stained.

## EXAMPLE 2: EXPRESSION OF RECOMBINANT TOLL PROTEIN IN BACTERIAL CELLS

In this example, TOLL protein is expressed as a recombinant glutathione-S-transferase (GST) fusion polypeptide in *E. coli* and the fusion polypeptide is isolated and characterized. Specifically, the TOLL sequence is fused to GST and this fusion polypeptide is expressed in *E. coli*, e.g., strain PEB199. Expression of the GST-TOLL fusion protein in PEB199 is induced with IPTG. The recombinant fusion polypeptide is purified from crude bacterial lysates of the induced PEB199 strain by affinity chromatography on glutathione beads. Using polyacrylamide gel electrophoretic analysis of the polypeptide purified from the bacterial lysates, the molecular weight of the resultant fusion polypeptide is determined.

**EXAMPLE 3:      EXPRESSION OF RECOMBINANT TOLL  
                     PROTEIN IN COS CELLS**

To express the TOLL gene in COS cells, the pcDNA/Amp vector by Invitrogen Corporation (San Diego, CA) is used. This vector contains an SV40 origin of replication, an ampicillin resistance gene, an *E. coli* replication origin, a CMV promoter followed by a polylinker region, and an SV40 intron and polyadenylation site. A DNA fragment encoding the entire TOLL protein and an HA tag (Wilson et al. (1984) *Cell* 37:767) or a FLAG tag fused in-frame to its 3' end of the fragment is cloned into the polylinker region of the vector, thereby placing the expression of the recombinant protein under the control of the CMV promoter.

To construct the plasmid, the TOLL DNA sequence is amplified by PCR using two primers. The 5' primer contains the restriction site of interest followed by approximately twenty nucleotides of the TOLL coding sequence starting from the initiation codon; the 3' end sequence contains complementary sequences to the other restriction site of interest, a translation stop codon, the HA tag or FLAG tag and the last 20 nucleotides of the TOLL coding sequence. The PCR amplified fragment and the pCDNA/Amp vector are digested with the appropriate restriction enzymes and the vector is dephosphorylated using the CIAP enzyme (New England Biolabs, Beverly, MA). Preferably the two restriction sites chosen are different so that the TOLL gene is inserted in the correct orientation. The ligation mixture is transformed into *E. coli* cells (strains HB101, DH5a, SURE, available from Stratagene Cloning Systems, La Jolla, CA, can be used), the transformed culture is plated on ampicillin media plates, and resistant colonies are selected. Plasmid DNA is isolated from transformants and examined by restriction analysis for the presence of the correct fragment.

COS cells are subsequently transfected with the TOLL-pcDNA/Amp plasmid DNA using the calcium phosphate or calcium chloride co-precipitation methods, DEAE-dextran-mediated transfection, lipofection, or electroporation. Other suitable methods for transfecting host cells can be found in Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 1989. The expression

of the TOLL polypeptide is detected by radiolabelling ( $^{35}\text{S}$ -methionine or  $^{35}\text{S}$ -cysteine available from NEN, Boston, MA. can be used) and immunoprecipitation (Harlow, E. and Lane, D. *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1988) using an HA specific monoclonal antibody. Briefly, the cells are labeled for 8 hours with  $^{35}\text{S}$ -methionine (or  $^{35}\text{S}$ -cysteine). The culture media are then collected and the cells are lysed using detergents (RIPA buffer, 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM Tris, pH 7.5). Both the cell lysate and the culture media are precipitated with an HA specific monoclonal antibody. Precipitated polypeptides are then analyzed by SDS-PAGE.

Alternatively, DNA containing the TOLL coding sequence is cloned directly into the polylinker of the pCDNA/Amp vector using the appropriate restriction sites. The resulting plasmid is transfected into COS cells in the manner described above, and the expression of the TOLL polypeptide is detected by radiolabelling and immunoprecipitation using a TOLL- specific monoclonal antibody.

#### Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

**Claims**

5

10

15

20

25

30

35

40

45

50

55

5

**What is claimed:**

10

1. An isolated nucleic acid molecule selected from the group consisting of:  
(a) a nucleic acid molecule comprising the nucleotide sequence set forth  
5 in SEQ ID NO:1;

15

(b) a nucleic acid molecule comprising the nucleotide sequence set forth  
in SEQ ID NO:3; and

(c) a nucleic acid molecule comprising the nucleotide sequence set forth  
in SEQ ID NO: 4.

10

20

2. An isolated nucleic acid molecule which encodes a polypeptide  
comprising the amino acid sequence set forth in SEQ ID NO: 2.

25

3. An isolated nucleic acid molecule comprising the nucleotide sequence  
15 contained in the plasmid deposited with ATCC® as Accession Number \_\_\_\_\_.

30

4. An isolated nucleic acid molecule which encodes a naturally occurring  
allelic variant of a polypeptide comprising the amino acid sequence set forth in SEQ ID  
NO: 2.

20

35

40

45

50

55



- 5                   5.     An isolated nucleic acid molecule selected from the group consisting of:
- 10                   a)     a nucleic acid molecule comprising a nucleotide sequence which  
is at least 60% homologous to the nucleotide sequence of SEQ ID NO:1, 3, or 4, or a  
complement thereof;
- 5                   b)     a nucleic acid molecule comprising a fragment of at least 491  
nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1 or 3,  
15                   or a complement thereof;
- c)     a nucleic acid molecule which encodes a polypeptide comprising  
an amino acid sequence at least about 60% homologous to the amino acid sequence of  
10     SEQ ID NO:2;
- 20                   d)     a nucleic acid molecule which encodes a fragment of a  
polypeptide comprising the amino acid sequence of SEQ ID NO:2, wherein the fragment  
comprises at least 15 contiguous amino acid residues of the amino acid sequence of SEQ  
25     ID NO:2; and
- 15                   e)     a nucleic acid molecule comprising a fragment of at least 15  
nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 4.
- 30                   6.     An isolated nucleic acid molecule which hybridizes to the nucleic acid  
molecule of any one of claims 1, 2, 3, 4, or 5 under stringent conditions.
- 20                   7.     An isolated nucleic acid molecule comprising a nucleotide sequence  
35                   which is complementary to the nucleotide sequence of the nucleic acid molecule of any  
one of claims 1, 2, 3, 4, or 5.
- 40                   8.     An isolated nucleic acid molecule comprising the nucleic acid molecule  
25                   of any one of claims 1, 2, 3, 4, or 5, and a nucleotide sequence encoding a heterologous  
polypeptide.
- 45                   9.     A vector comprising the nucleic acid molecule of any one of claims 1, 2,  
30     3, 4, or 5.
- 50
- 55

5

10. The vector of claim 9, which is an expression vector.

10

11. A host cell transfected with the expression vector of claim 10.

5

12. A method of producing a polypeptide comprising culturing the host cell of claim 11 in an appropriate culture medium to, thereby, produce the polypeptide.

15

13. An isolated polypeptide selected from the group consisting of:

20

a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2;

25

b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule consisting of SEQ ID NO:1, 3, or 4 under stringent conditions;

30

c) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 60% homologous to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, 3, or 4;

35

d) a polypeptide comprising an amino acid sequence which is at least 60% homologous to the amino acid sequence of SEQ ID NO:2.

40

14. The isolated polypeptide of claim 13 comprising the amino acid sequence of SEQ ID NO:2.

25

15. The polypeptide of claim 13, further comprising heterologous amino acid sequences.

45

16. An antibody which selectively binds to a polypeptide of claim 13.

50

55

5

17. A method for detecting the presence of a polypeptide of claim 13 in a sample comprising:

10

a) contacting the sample with a compound which selectively binds to the polypeptide; and

5

b) determining whether the compound binds to the polypeptide in the sample to thereby detect the presence of a polypeptide of claim 13 in the sample.

15

18. The method of claim 17, wherein the compound which binds to the polypeptide is an antibody.

10

20

19. A kit comprising a compound which selectively binds to a polypeptide of claim 13 and instructions for use.

25

20. A method for detecting the presence of a nucleic acid molecule of any one of claims 1, 2, 3, 4, or 5 in a sample comprising:

30

a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and

35

b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample to thereby detect the presence of a nucleic acid molecule of any one of claims 1, 2, 3, 4, or 5 in the sample.

40

21. The method of claim 20, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

45

50

22. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of any one of claims 1, 2, 3, 4, or 5 and instructions for use.

55

50

55

5  
23. A method for identifying a compound which binds to a polypeptide of claim 13 comprising:

- 10 a) contacting the polypeptide, or a cell expressing the polypeptide with a test compound; and  
5 b) determining whether the polypeptide binds to the test compound.

15 24. The method of claim 23, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

- 10 a) detection of binding by direct detection of test compound/polypeptide binding;  
20 b) detection of binding using a competition binding assay; and  
c) detection of binding using an assay for TOLL activity.

25 25. A method for modulating the activity of a polypeptide of claim 13 comprising contacting the polypeptide or a cell expressing the polypeptide with a  
15 compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.  
30

26. A method for identifying a compound which modulates the activity of a  
20 polypeptide of claim 13 comprising:

- 35 a) contacting a polypeptide of claim 13 with a test compound; and  
b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the  
40 polypeptide.

1/6

## Human TOLL cDNA sequence:

GTCTGACCCACGCGTCCGCGGACGCGTGGGGTTGGATTTTTTCAAAGAGTAAA  
CCAGACCCGTTGACCAAGGTGTAGACTAAGAAGTGGAGTCATGCTTCACACGG  
CCATATCATGCTGGCAGCCATTCTGGGTCTGGCTGTGGTGTAAATCTTCATG  
GGATCCACCATTTGGCTGCCCCGCTCGCTGTGAGTGCTCTGCCCAGAACAAATC  
TGTTAGCTGTACAGAAGCGATTGATCGCCATCCCAGAGGGCATTCCCCTC  
GAAACCAAATCTTGGACCTCAGTAAAAACAGGCTAAAAAGCGTCAACCCCTG  
AAGAATTATATCATATCCTCTGCTGGAAGAGATAGACTTGAGTGACAACAT  
CATTGCCAATGTGGAACAGGAGCATTCAACAATCTCTTTAACCTGCGTTCCC  
TCCGCCTAAAAGGCAATCGTCTAAAGCTGGTCCCTTTGGGAGTATTCACGGG  
GCTGTCCAATCTCACTAAGCTTGACATTAGTGAGAATAAGATTGTCATTTTAC  
TAGACTACATGTTCCAAGATCTACATAACCTGAAGTCTCTAGAAGTGGGGGA  
CAATGATTTGGTTTATATATCACACAGGGCATTCAAGTGGGCTTCTTAGCTTG  
AGCAGCTCACCTTGAGAAATGCAACTTAACAGCAGTACCAACAGAAGCCCT  
CTCCCACTCCGCGAGCCTCATCAGCCTGCATCTGAAGCATCTCAATATCAACA  
ATATGCCTGTGTATGCCTTTAAAAGATTGTTCCACCTGAAACACCTAGAGATT  
GACTATTGGCCCTTACTGGATATGATGCCTGCCAATAGCCTCTACGGTCTCAA  
CCTCACATCCCTTTCAGTCACCAACACCAATCTGTCTACTGTACCCTTCTCTTG  
CTTTAAACACCTGGTATACCTGACTCACCTTAACCTCTCCTACAATCCCATCA  
GCACTATTGAAGCAGGCATGTTCTCTGACCTGATCCGCCTTCAGGAGCTTCAT  
ATAGTGGGGGCCAGCTTCGCACCATTTAGCCTCACTCCTTCCAAGGGCTCCG  
CTTCTACGCGTGTCAATGTGTCTCAGAACCCTGCTGGAACTTTGGAAGAGA  
ATGTCTTCTCCTCCCTAGGGCTCTGGAGGTCTTGAGCATTAACAACAACCTT  
CTGGCCTGTGACTGCCGCTTCTCTGGATCTTGACGCGACAGCCACCCTGCA  
GTTTGGTGGCCAGCAACCTATGTGTGCTGGCCAGACACCATCCGTGAGAGG  
TCTTTCAAGGATTTCCATAGCACTGCCCTTTCTTTTACTTTTACCTGCAAAAAA  
CCCAAAATCCGCTTTGCCCAGGATCAAGACAGCGGGATGTATGTTTGCATCG  
CTAGCAATGCTGTGGGAATGATACCTTCACAGCCTCCTTAAGTGTGAAAGG  
ATTGCGTTTCAATCGTTTTCTTTATGCGAACAGGACCCCTATGTACATGACCG  
ACTCCAATGACACCATTTCATGGCACCATGCAATACCTTTTCCCTGGAC  
CTTAAACAATACTGGTGTCTACAGCTATGGGCTGCTTCACATTCCTGGGAGT  
GGTTTTATTTTGTCTTCTCTCTTTTGTGTGGAGCCGAGGGAAAGGCAAGC  
ACAAAAACAGCATTGACCTTGAGTATGTGCCGAAAAAACAATGGTGTGT  
TGTGGAAGGGGAGGTAGCTGGACCCAGGAGGTTCAACATGAAATGATTGTA  
AGGCCCACCCCTCACATTACTGTCTCTTTGTCAATGTGGGTAAATCAGTAAGAC  
AGTATGGCACAGTAAATTACTAGATTAAGAGGCAGCCATGTGCAGCTGCCCC  
TGTATCAAAAGCAGGGTCTATGGAAGCAGGAGGACTTCAATGGAGACTCTC  
CATCGAAAGGCAGGCAGGCAGGCATGTGTGAGAGCCCTTCACACAGTGGGAT  
ACTAAGTGTGTGCGTTGCAATATTGGCGTTCTGGGGATCTCAGTAATGAACC  
TGAATATTTGGCTCACACTCACGGACAATTATTCAGCATTTTCTACCACTGCA  
AA  
AA

Fig. 1

SUBSTITUTE SHEET (RULE 26)

2/6

Human TOLL coding region:

ATGCTTCACACGGCCATATCATGCTGGCAGCCATTCTGGGTCTGGCTGTGGT  
GTTAATCTTCATGGGATCCACCATTGGCTGCCCCGCTCGCTGTGAGTGTCTG  
CCCAGAACAATCTGTTAGCTGTCACAGAAGGCGATTGATCGCCATCCCAGA  
GGGCATTCCCATCGAAACCAAATCTTGGACCTCAGTAAAAACAGGCTAAAA  
AGCGTCAACCCTGAAGAATTTCATATCATATCCTCTGCTGGAAGAGATAGACT  
TGAGTGACAACATCATTGCCAATGTGGAACCAGGAGCATTCAACAATCTCTT  
TAACCTGCGTTCCCTCCGCCTAAAAGGCAATCGTCTAAAGCTGGTCCCTTTGG  
GAGTATTCACGGGGCTGTCCAATCTCACTAAGCTTGACATTAGTGAGAATAA  
GATTGTCAATTTACTAGACTACATGTTCCAAGATCTACATAACCTGAAGTCTC  
TAGAAGTGGGGGACAATGATTTGGTTTATATATCACACAGGGCATTCACTGG  
GCTTCTTAGCTTGGAGCAGCTCACCTGGAGAAATGCAACTTAACAGCAGTA  
CCAACAGAAGCCCTCTCCACCTCCGAGCCTCATCAGCCTGCATCTGAAGC  
ATCTCAATATCAACAATATGCCTGTGTATGCCTTTAAAAGATTGTTCCACCTG  
AAACACCTAGAGATTGACTATTGGCCTTTACTGGATATGATGCCCTGCCAATAG  
CCTCTACGGTCTCAACCTCACATCCCTTTCAGTCACCAACACCAATCTGTCTA  
CTGTACCCTTCCCTTGCTTTAAACACCTGGTATACCTGACTCACCTTAACCTCT  
CCTACAATCCCATCAGCACTATTGAAGCAGGCATGTTCTCTGACCTGATCCGC  
CTTCAGGAGCTTCATATAGTGGGGGCCAGCTTCGCACCATTGAGCCTCACTC  
CTTCCAAGGGCTCCGCTTCCCTACGCGTGCTCAATGTGTCTCAGAACCCTGCTGG  
AACTTTGGAAGAGAATGTCTTCTCCTCCCTAGGGCTCTGGAGGTCTTGAGC  
ATTAACAACAACCCCTCTGGCCTGTGACTGCCGCCCTTCTCTGGATCTTGCAGCG  
ACAGCCCACCCTGCAGTTTGGTGGCCAGCAACCTATGTGTGCTGGCCCAGAC  
ACCATCCGTGAGAGGTCTTTCAAGGATTTCCATAGCACTGCCCTTTCTTTTTTA  
CTTTACCTGCAAAAAACCCAAAATCCGCTTTGCCCAGGATCAAGACAGCGGG  
ATGTATGTTTGCATCGCTAGCAATGCTGCTGGGAATGATACCTTCACAGCCTC  
CTTAACGTGTGAAAGGATTTCGCTTCAGATCGTTTCTTTATGCGAACAGGACCC  
CTATGTACATGACCGACTCCAATGACACCATTTCCAATGGCACCAATGCCAAT  
ACTTTTCCCTGGACCTTAAAACAATACTGGTGTCTACAGCTATGGGCTGCTT  
CACATTCCTGGGAGTGGTTTTATTGTTTTCTCTCCTTTTTGTGTGGAGCCG  
AGGGAAAGGCAAGCACAAAACAGCATTGACCTTGAGTATGTGCCAGAAA  
AAACAATGGTGTGTGTGGAAGGGGAGGTAGCTGGACCCAGGAGGTTCAAC  
ATGAAAATGATTGA

Fig. 1 (continued)

3/6

Human TOLL protein sequence:

MLHTAISCWQPFGLAVVLIFMGSTIGCPARCECSAQNKSVSCHRRRLIAIPEGIPI  
ETKIIDL SKNRLKSVNPEEFISYPLLEEIDLSDNIIANVEPGA FN NLFNLRSLRLKG  
NRLKLVPLGVFTGLSNLTKLDISENKIVILLDYMFDLHNLKSLEVGDNDLVYIS  
HRAFSGLLSLEQLTEKCNLTAVPTEALSHLRSLISLHLKHLNINMPVYAFKRLF  
HLKHLIDYWPLLDMPANSYGLNLTSLSVTNTNLSTVPFLAFKHLVYLTHLN  
LSYNPISTIEAGMFSDLIRLQELHIVGAQLRTIEPHSFQGLRFLRVLNVSQNLLLETL  
EENVFSSPRALEVLSINNNPLACDCRLLWILQRQPTLQFGGQQPMCAGPDTIRERS  
FKDFHSTALSFYFTCKKPKIRFAQDQDSGMYVCIASNAAGNDTFTASLTVKGFAS  
DRFLYANRTPMYMTDSNDTISNGTNANTFSLDLKTILVSTAMGCFTFLGVVLFCE  
LLLFVWSRGKGKHKNSIDLEYVPRKNGAVVEGEVAGPRRFNMKMI.

Fig. 1 (continued)

4/6

HMMER 2.1.1 (DEC 1998)

Copyright (C) 1992-1998 Washington University School of Medicine

HMMER is freely distributed under the GNU General Public License (GPL)

HMM file: /prod/ddm/seqanal/PFAM/pfam3.4/Pfam

Sequence file: /tmp/orfanal.28310.aa

Query: mill-toll

Scores for sequence family classification (score includes all domains):

Model	Description	Score	E-value	N
LRR	PF00560 Leucine Rich Repeat (2 copies)	183.5	3.3e-51	6

Parsed for domains:

Model	Domain	seq-f	seq-t	hmm-f	hmm-t	score	E-value
LRR	1/6	58	105	1	48	32.3	1.1e-05
LRR	2/6	106	153	1	48	43.9	3.7e-09
LRR	3/6	154	201	1	48	37.6	2.8e-07
LRR	4/6	202	250	1	48	17.1	0.42
LRR	5/6	251	297	1	48	34.9	1.8e-06
LRR	6/6	298	345	1	48	47.4	3.1e-10

Alignments of top-scoring domains:

LRR: domain 1 of 6, from 58 to 105: score 32.3, E = 1.1e-05

```

*->nLeeLdLsnNkLtslppgalsnLpnLeeLdLsnNnLtslppglfqnl
+ LdLs N+L+s+ p+ + + p Lee+dLs+N + ++pg+f nL
mill-toll 58 ETKILDLSKNRLKSVNPEEPISYPLLEEIDLSNIIANVEPGAFFNL 104
k<-*
```

mill-toll 105 F 105

LRR: domain 2 of 6, from 106 to 153: score 43.9, E = 3.7e-09

```

*->nLeeLdLsnNkLtslppgalsnLpnLeeLdLsnNnLtslppglfqnl
nL++L+L +N+L+ +p g++ +L+nL++Ld+s N++ 1 ++ fq+L
mill-toll 106 NLRSLRLKGNRLKLVPLGVFTGLSNLTKLDISENKIVILLDYMFQDL 152
k<-*
```

mill-toll 153 H 153

LRR: domain 3 of 6, from 154 to 201: score 37.6, E = 2.8e-07

```

*->nLeeLdLsnNkLtslppgalsnLpnLeeLdLsnNnLtslppglfqnl
nL++L++ +N L ++ a+s+L +Le L L +nLt +p +++++L
mill-toll 154 NLKSLEVGDNLDVYISHRAFSGLLSLEQLTEKCNLTAVPTEALSHL 200
k<-*
```

mill-toll 201 R 201

Fig. 2

SUBSTITUTE SHEET (RULE 26)



5/6

LRR: domain 4 of 6, from 202 to 250: score 17.1, E = 0.42

\*->nLeeLdLsnNkLtslppgalsnLpnLeeLdLsnNnLt.slppglfq  
 +L +L+L + ++ + p a++ L +L++L +++ L + p +++ +  
 mill-toll 202 SLISLHLKHLNINNMVYAFKRLFHLKHLLEIDYWPLLDMPANSLYG 248

Lk&lt;-\*

L+

mill-toll 249 LN 250

LRR: domain 5 of 6, from 251 to 297: score 34.9, E = 1.8e-06

\*->nLeeLdLsnNkLtslppgalsnLpnLeeLdLsnNnLtslppglfqL  
 L++L+++n++L+++p a+++L L++L+Ls+N +++++ g f++L  
 mill-toll 251 -LTSLSVTETNLSTVPFLAFKHLVYLTHLNLSYNPISTIEAGMPSDL 296

k&lt;-\*

mill-toll 297 I 297

LRR: domain 6 of 6, from 298 to 345: score 47.4, E = 3.1e-10

\*->nLeeLdLsnNkLtslppgalsnLpnLeeLdLsnNnLtslppglfqL  
 +L+eL++ + +L++++p+++++L+ L++L++s N L++l++++f++  
 mill-toll 298 RLQELHIVGAQLRTIEPHSFQGLRFLRVLNVSQNLEETLEENVFSSP 344

k&lt;-\*

+

mill-toll 345 R 345

//

Fig. 2 (continued)

6/6

MOUSE TOLL DNA

ACTGTCCTCTGTTACTGTTGGTCGTGAGTAAGACGCTCTGATAGAGTGACTCG  
ATCACAAGGTTATCGGGCAGCTTTGCGCAGCTGCCCTGTGTCAAAGCAGGG  
TCCATGGAAGCAGGAAGACTTCTCATGGAGACTGGCTGATTAGAGGCAGGCA  
GGCATGTGTCAGAGCCCTTCACACAGTGGGATACTAATTGTTTGCATTGCAAA  
TATTGGCATTCTGGGGATCTCAGCAATGAACCTGAACCTTTGGCTCATGCTGA  
TGGACAATAATTCAACATTTTCTACCCTGCAAACTAAAAGGAAAAAAT  
TAAAAGAACAACCTACAGTGTAGGATTTACATATTAAAAGACACATTTGT  
CTAAACATACTCTACAGTCAAATTTGTATTTATTATCATTTGTTAAACCTT  
GCATCATACAATACTGTTGGTTCAGCACCAAAAAGAGATCAATATATCTTTT  
TTTTGAAACATATATGCTGTATATGTTTAAAGCAATATGAATGAGAGGTTGT  
GCTTTTAGTTACTCACCAGTATAGATCCAAGTGTGGTTTCACCTTCCTTTTACC  
TGCAGATAAACCTGAGAATAGATCCCTGGAATACTAGGCAGAGATGTGTTGA  
GATGTGTWTGTCTGATGTAGGATGCCAAGAAACAAGAMCCAAGTCMAAACT  
GCTCMACTCTGTAACTTCTGTTACTATAAATAAAGGCATGTGCCTAGTTTGT  
ATACAAAAAARGCGGC

Fig. 3